arXiv:0909.1730v1 [math.QA] 9 Sep 2009

Two-parameter quantum vertex representations via finite groups and the McKay correspondence

Naihuan Jing and Honglian Zhang*

Abstract. We introduce two-parameter quantum toroidal algebras of simply laced types and provide their group theoretic realization using finite subgroups of $SL_2(\mathbb{C})$ via McKay correspondence. In particular our construction contains a realization of the vertex representation of the two-parameter quantum affine algebras of ADE types.

1. Introduction

In a series papers [**FJW1, FJW2**] the basic representations of two-toroidal Lie algebras and their quantum analogs, including affine Lie algebras and quantum affine algebras of simply laced types as subaglebras, were constructed from representation theory of finite groups of $SL_2(\mathbb{C})$ via the celebrated McKay correspondence. Using purely representation theoretic data the Frenkel-Kac [**FK**] and Frenkel-Jing [**FJ**] vertex representations are constructed as a by-product of the unified group theoretic constructions from the root lattice of the corresponding finite dimensional Lie algebra $\mathfrak{g}$. In [**J4**] we have pointed out that such a uniformed construction incorporate not only the toroidal Lie algebras, quantum toroidal algebras [**VV**] (see also [**J3**]) but also give a general algebraic machinery to realize other related algebraic structures.

In the current paper we provide a two-parameter quantum analog of the toroidal Lie algebras of simply laced types using the new form of McKay correspondence. In particular this gives a group theoretic realization of the newly revitalized two-parameter quantum affine algebras [**HRZ, Z**] as distinguished subalgebras of our new quantum toroidal algebras. As expected, our construction degenerates to the quantum affine case by specializing $r = s^{-1}$. Through this new construction we have also shown that the vertex representation constructed in [**HZ, Z**] is a natural generalization of Frenkel-Jing construction and also reconfirm the two-parameter generalization of the usual quantum affine algebras and their Drinfeld realization. Our construction further shows that the toroidal version is more suitable and natural in this picture and reveal more symmetry in the structure of the two parameter quantum toroidal algebras.

2000 *Mathematics Subject Classification.* Primary 17B37, 17B67, 17B69, 81R10.

Key words and phrases. Two-parameter quantum toroidal algebra, finite groups, wreath products, McKay correspondence.

*H.Z., Corresponding Author.

The new two-parameter quantum toroidal Lie algebras or double affine algebras provide a new layer of generalization or quantization. To show the relationships we can depict them in the following diagram similar to that given by I. Frenkel earlier. Our new algebra adds a new direction of generalization.

$$\begin{tikzcd}
 & & \mathfrak{g} \arrow[dl] \arrow[dr] & & \\
 & \widehat{\mathfrak{g}} \arrow[dl] \arrow[dr] & & U_q(\mathfrak{g}) \arrow[dl] \arrow[dr] & \\
\widehat{\widehat{\mathfrak{g}}} \arrow[dr] & & U_q(\widehat{\mathfrak{g}}) \arrow[dl] \arrow[dr] & & U_{r,s}(\mathfrak{g}) \arrow[dl] \\
 & U_q(\widehat{\widehat{\mathfrak{g}}}) \arrow[dr] & & U_{r,s}(\widehat{\mathfrak{g}}) \arrow[dl] & \\
 & & U_{r,s}(\widehat{\widehat{\mathfrak{g}}}) & &
\end{tikzcd}$$

It is amazing that each vertex in the diagram admits a realization through McKay correspondence, and each level of complexity is achieved by replacing the finite group Γ by $\Gamma \times \mathbb{C}^\times$ and by $\Gamma \times \mathbb{C}^\times \times \mathbb{C}^\times$. The quantum parameter q and (r, s) are respectively represented by special characters on the group $\mathbb{C}^\times$ and $\mathbb{C}^\times \times \mathbb{C}^\times$.

At the early stage of the development of quantum groups it was already realized there exist multi-parameter quantum groups [**R, T**]. Though two-parameter cases play an important role in the dual quantum picture, the development of two-parameter quantum groups only took a turn after more structures are revealed in series of papers of Benkart and Witherspoon [**BW1, BW2, BW3**] for type A. Shortly after these generalizations to other types are given by Bergeron, Gao and Hu [**BGH1, BGH2, BH, HS**]. Recently Hu, Rosso and one of us found the two parameter quantum affine algebras of ADE types [**HRZ, Z**] via their vertex representations [**HZ**] based on generalized Drinfeld realizations (cf. [**J2**]), which are two-parameter analog of the Frenkel-Jing representations [**FJ**]. In all these work one realizes that the two-parameter analog, like its one-parameter case, amounts to a clever deformation of the natural number n to the two-parameter quantum number:

$$[n] = \frac{r^n - s^n}{r - s} = r^{n-1} + r^{n-2}s + \cdots + rs^{n-2} + s^{n-1}. \tag{1.1}$$

Clearly when $rs = 1$, the two-parameter quantum number degenerates to the one-parameter quantum number.

As in the previous construction of McKay correspondence and quantum toroidal algebras we recover the basic representation of $U_{r,s}(\widehat{\mathfrak{g}})$ by choosing

$$\xi = \gamma_0 \otimes ((rs^{-1})^{\frac{1}{2}} + (r^{-1}s)^{\frac{1}{2}}) - \pi \otimes 1_{\mathbb{C}^\times \times \mathbb{C}^\times},$$

where γ_0, $1_{\mathbb{C}^\times \times \mathbb{C}^\times}$ are the trivial characters of Γ and $\mathbb{C}^\times \times \mathbb{C}^\times$ respectively, r and s are two independent natural characters of $\mathbb{C}^\times$, and π is the natural character of the imbedding of Γ in $SL_2(\mathbb{C})$. The natural appearance of the two parameter quantum toroidal algebra in this picture intrinsically shows its importance in representation theory of two parameter quantum affine algebras.

The paper is organized as follows. Section 2 recalls the basic material of wreath products Γ_n of symmetric groups associated with any finite group and the Hopf algebra structures in the representation ring. Section 3 studies the representation ring $R(\Gamma_n \times \mathbb{C}^\times \times \mathbb{C}^\times)$ and weighted bilinear forms. Section 4 gives the two-parameter McKay weights for each finite subgroup of $SL_2(\mathbb{C})$. Section 5 defines the two-parameter Heisenberg algebra and realizes its canonical representation using group theoretic data out of Γ. Section 6 gives the Frobenius-type characteristic map between $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ and $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$. Section 7 realizes the two-parameter quantum vertex operators using irreducible characters of $\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times$, and finally in Section 8 we introduce two-parameter quantum toroidal algebras and provide their realization via McKay correspondence, and in particular this also provides a group theoretic realization of the basic representation of the two-parameter quantum affine algebras.

2. Wreath products and vertex representations

2.1. The wreath product Γ_n. Let Γ be a finite group and n a non-negative integer. The wreath product Γ_n is the semidirect product of the n-th direct product $\Gamma^n = \Gamma \times \cdots \times \Gamma$ and the symmetric group S_n:

$$\Gamma_n = \{(g, \sigma) | g = (g_1, \dots, g_n) \in \Gamma^n, \sigma \in S_n\}$$

with the group multiplication

$$(g, \sigma) \cdot (h, \tau) = (g\,\sigma(h), \sigma\tau),$$

where S_n acts on Γ^n by permuting the factors.

Let Γ_* be the set of conjugacy classes of Γ consisting of $c^0 = \{1\}$, c^1, $\dots$, c^ℓ and Γ^* be the set of $\ell + 1$ irreducible characters: $\gamma_0, \gamma_1, \dots, \gamma_\ell$, where γ_0 is the trivial character of Γ. The order of the centralizer of an element in the conjugacy class c is denoted by ζ_c, so the order of the conjugacy class c is $|c| = |\Gamma|/\zeta_c$, where $|\Gamma|$ is the order of Γ.

A partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$ is a decomposition of $n = |\lambda| = \lambda_1 + \cdots + \lambda_l$ with nonnegative integers: $\lambda_1 \geq \cdots \geq \lambda_\ell \geq 1$, where $\ell = \ell(\lambda)$ is called the *length* of the partition λ and λ_i are called the *parts* of λ. Another notation for λ is

$$\lambda = (1^{m_1} 2^{m_2} \cdots)$$

with m_i being the multiplicity of parts equal to i in λ. Denote by $\mathcal{P}$ the set of all partitions of integers and by $\mathcal{P}(S)$ the set of all partition-valued functions on a set S. The weight of a partition-valued function $\rho = (\rho(s))_{s\in S}$ is defined to be $\|\rho\| = \sum_{s\in S} |\rho(s)|$. We also denote by $\mathcal{P}_n$ (resp. $\mathcal{P}_n(S)$) the subset of $\mathcal{P}$ (resp. $\mathcal{P}(S)$) of partitions with weight n.

It is well-known that the conjugacy classes of Γ_n are parameterized by partition-valued functions on Γ_*. Let $x = (g, \sigma) \in \Gamma_n$, where $g = (g_1, \dots, g_n) \in \Gamma^n$ and $\sigma \in S_n$ is presented as a product of disjoint cycles. For each cycle $(i_1 i_2 \cdots i_k)$ of σ, we define the *cycle-product* element $g_{i_k} g_{i_{k-1}} \cdots g_{i_1} \in \Gamma$, which is determined up to conjugacy in Γ by g and the cycle. For any conjugacy class $c \in \Gamma$ and each integer $i \geq 1$, the number of i-cycles in σ whose cycle-product lies in c will be denoted by $m_i(c)$. This gives rise to a partition $\rho(c) = (1^{m_1(c)} 2^{m_2(c)} \dots)$ for $c \in \Gamma_*$. Thus we obtain a partition-valued function $\rho = (\rho(c))_{c\in\Gamma_*} \in \mathcal{P}(\Gamma_*)$ such that $\|\rho\| = \sum_{i,c} im_i(\rho(c)) = n$. This is called the *type* of the element (g, σ). It is known [**M**] that two elements in the same conjugacy class have the same type and there exists a one-to-one correspondence between the sets $(\Gamma_n)_*$ and $\mathcal{P}_n(\Gamma_*)$. We will say that ρ is the type of the conjugacy class of Γ_n.

Given a class c we denote by c^{-1} the class $\{x^{-1} | x \in c\}$. For each $\rho \in \mathcal{P}(\Gamma_*)$ we also associate the partition-valued function

$$\overline{\rho} = (\rho(c^{-1}))_{c\in\Gamma_*}.$$

Given a partition $\lambda = (1^{m_1}2^{m_2}\ldots)$, we denote by $z_\lambda = \prod_{i\geq 1} i^{m_i} m_i!$ the order of the centralizer of an element of cycle type λ in $S_{|\lambda|}$. The order of the centralizer of an element $x = (g, \sigma) \in \Gamma_n$ of type $\rho = (\rho(c))_{c\in\Gamma_*}$ is given by

$$Z_\rho = \prod_{c\in\Gamma_*} z_{\rho(c)} \zeta_c^{l(\rho(c))}.$$

2.2. Grothendieck ring $R_{\Gamma\times C^\times\times C^\times}$.

Let $R_{\mathbb{Z}}(\Gamma)$ be the $\mathbb{Z}$-lattice generated by γ_i, $i = 0, \ldots, r$, and $R(\Gamma) = \mathbb{C} \otimes R_{\mathbb{Z}}(\Gamma)$ be the space of complex class functions on the group Γ. In the previous work on the McKay correspondence and vertex representations [**W, FJW1**], the Grothendieck ring $R_\Gamma = \bigoplus_{n\geq 0} R(\Gamma_n)$ was studied. In quantum case, the Grothendieck ring was $R_{\Gamma\times\mathbb{C}^\times} = \bigoplus_{n\geq 0} R(\Gamma_n \times \mathbb{C}^\times)$ [**FJW2**]. In our two-parameter quantum case we need to add the another ring $R(\mathbb{C}^\times)$, the space of characters of $\mathbb{C}^\times \times \mathbb{C}^\times = \{(t_1, t_2) \in \mathbb{C} \times \mathbb{C} | t_1, t_2 \neq 0\}$.

Let r, s be the irreducible character of $\mathbb{C}^\times$ that sends t to itself. Then $R(\mathbb{C}^\times \times \mathbb{C}^\times)$ is spanned by irreducible multiplicative characters $r^m s^n$, m, $n \in \mathbb{Z}$, where

$$r^m(t_1) = t_1^m, \qquad s^n(t_2) = t_2^n \qquad t_1, t_2 \in \mathbb{C}^\times.$$

Thus $R(\mathbb{C}^\times \times \mathbb{C}^\times)$ is identified with the ring $\mathbb{C}[r^{\pm 1}, s^{\pm 1}]$, and we have

$$R(\Gamma \times \mathbb{C}^\times \times \mathbb{C}^\times) = R(\Gamma) \otimes R(\mathbb{C}^\times \times \mathbb{C}^\times).$$

An elements of $R(\Gamma \times \mathbb{C}^\times \times \mathbb{C}^\times)$ can be written as a finite sum:

$$f = \sum_i f_i \otimes r^{m_i} s^{n_i}, \qquad f_i \in R(\Gamma), m_i, n_i \in \mathbb{Z}.$$

We can also view f as a function on Γ with values in the ring of Laurent polynomials $\mathbb{C}[r^{\pm 1}, s^{\pm 1}]$. In this case we will write $f^{r,s}$ to indicate the formal variables r, s, then $f^{r,s}(c) = \sum_i f_i(c) r^{m_i} s^{n_i} \in \mathbb{C}[r^{\pm 1}, s^{\pm 1}]$. As a function on $\Gamma \times \mathbb{C}^\times \times \mathbb{C}^\times$, we have $f(c, t_1, t_2) = \sum_i f_i(c) t_1^{m_i} t_2^{n_i}$.

Denote by $R_{\Gamma\times C^\times\times C^\times}$ the following direct sum:

$$R_{\Gamma\times C^\times\times C^\times} = \bigoplus_{n\geq 0} R(\Gamma_n \times \mathbb{C}^\times \times \mathbb{C}^\times) \simeq R_\Gamma \otimes \mathbb{C}[r^{\pm 1}, s^{\pm 1}].$$

2.3. Hopf algebra structure on $R_{\Gamma\times C^\times\times C^\times}$.

The multiplication m in $\mathbb{C}^\times \times \mathbb{C}^\times$ and the diagonal map $\mathbb{C}^\times \times \mathbb{C}^\times \xrightarrow{d} \mathbb{C}^\times \times \mathbb{C}^\times \times \mathbb{C}^\times \times \mathbb{C}^\times$ induce the Hopf algebra structure on $R(\mathbb{C}^\times \times \mathbb{C}^\times)$.

(2.1)

$$m_{\mathbb{C}^\times\times C^\times} : R(\mathbb{C}^\times \times \mathbb{C}^\times) \otimes R(\mathbb{C}^\times \times \mathbb{C}^\times) \xrightarrow{\cong} R(\mathbb{C}^\times \times \mathbb{C}^\times \times \mathbb{C}^\times \times \mathbb{C}^\times) \xrightarrow{d^*} R(\mathbb{C}^\times \times \mathbb{C}^\times),$$

(2.2)

$$\Delta_{\mathbb{C}^\times\times C^\times} : R(\mathbb{C}^\times) \xrightarrow{m^*} R(\mathbb{C}^\times \times \mathbb{C}^\times) \xrightarrow{\cong} R(\mathbb{C}^\times) \otimes R(\mathbb{C}^\times).$$

In terms of the basis $\{r^m s^n\}$ we have

$$
\begin{aligned}
r^{m_1}s^{n_1}\cdot r^{m_2}s^{n_2} &= r^{m_1+m_2}s^{n_1+n_2},\\
\Delta(r^m s^n) &= r^m s^n \otimes r^m s^n,
\end{aligned}
$$

where we abbreviate $\Delta_{\mathbb{C}^\times\times\mathbb{C}^\times}$ by Δ and follow the convention of writing $a\cdot b = m_{\mathbb{C}^\times\times\mathbb{C}^\times}(a\otimes b)$.

The antipode $S_{\mathbb{C}^\times\times\mathbb{C}^\times}$ and the counit $\epsilon_{\mathbb{C}^\times\times\mathbb{C}^\times}$ are given by

$$
S_{\mathbb{C}^\times\times\mathbb{C}^\times}(r^m s^n) = r^{-m}s^{-n}, \qquad \epsilon_{\mathbb{C}^\times\times\mathbb{C}^\times}(r^m s^n) = \delta_{m-n,0}.
$$

We extend the Hopf algebra structures on $R(\mathbb{C}^\times\times\mathbb{C}^\times)$ and R_Γ into a Hopf algebra structure on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ using a standard procedure in Hopf algebra [**A**]. The multiplication and comultiplication are given by the respective composition of the following maps:

$$
\begin{aligned}
m: R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)\otimes R(\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times) &\xrightarrow{\cong} R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times\times\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times)\\
&\xrightarrow{1\otimes m_{\mathbb{C}^\times\times\mathbb{C}^\times}} R(\Gamma_n\times\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times)\xrightarrow{Ind\otimes 1} R(\Gamma_{n+m}\times\mathbb{C}^\times\times\mathbb{C}^\times); \qquad (2.3)
\end{aligned}
$$

$$
\begin{aligned}
\Delta: R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times) &\xrightarrow{Res\otimes 1} \oplus_{m=0}^{n} R(\Gamma_{n-m}\times\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times)\\
&\xrightarrow{1\otimes\Delta_{\mathbb{C}^\times\times\mathbb{C}^\times}} \oplus_{m=0}^{n} R(\Gamma_{n-m}\times\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times\times\mathbb{C}^\times\times\mathbb{C}^\times)\\
&\xrightarrow{\cong} \oplus_{m=0}^{n} R(\Gamma_{n-m}\times\mathbb{C}^\times\times\mathbb{C}^\times)\otimes R(\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times), \qquad (2.4)
\end{aligned}
$$

where we have used the identification of $R(\mathbb{C}^\times\times\mathbb{C}^\times)$ with $R(\mathbb{C}^\times)\otimes R(\mathbb{C}^\times)$ in (2.1-2.2). Also $Ind: R(\Gamma_n\times\Gamma_m)\longrightarrow R(\Gamma_{n+m})$ denotes the induction functor and $Res: R(\Gamma_n)\longrightarrow R(\Gamma_{n-m}\times\Gamma_m)$ denotes the restriction functor.

The antipode is given by

$$
S(f(g,(t_1,t_2))) = f(g^{-1},(t_1^{-1},t_2^{-1})), \qquad g\in\Gamma, t\in\mathbb{C}^\times\times\mathbb{C}^\times.
$$

In particular, $S(\gamma)(c)=\gamma(c^{-1})$ for $\gamma\in\Gamma^*$. As we mentioned earlier, we may write $f\in R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ as

$$
f^{r,\,s}(g) = \sum_i f_i(g) r^{m_i} s^{n_i},
$$

Then $S(f^{r,\,s})(g) = \sum_i f_i(g^{-1}) r^{-m_i}s^{-n_i}$.

The counit ϵ is defined by

$$
\epsilon(R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)) = 0, \qquad \text{if} \quad n\neq 0,
$$

and ϵ on $R(\mathbb{C}^\times\times\mathbb{C}^\times)$ is the counit of the Hopf algebra $R(\mathbb{C}^\times\times\mathbb{C}^\times)$.

3. A weighted bilinear form on $R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)$

3.1. A standard bilinear form on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$. Let $f,g\in R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ with $f=\sum_i f_i\otimes r^{m_i}s^{n_i}$ and $g=\sum_i g_i\otimes r^{k_i}s^{l_i}$. The $\mathbb{C}[r^{\pm1},s^{\pm1}]$-valued standard $\mathbb{C}$-bilinear form on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ is defined as

$$
\begin{aligned}
\langle f,g\rangle_\Gamma^{r\,s} &= \sum_{i,j}\langle f_i,g_j\rangle_\Gamma r^{m_i-k_j}s^{n_i-l_j}\\
&= \sum_{i,j}\sum_{c\in\Gamma_*}\zeta_c^{-1}f_i(c)g_j(c^{-1})r^{m_i-k_j}s^{n_i-l_j},
\end{aligned}
$$

where we recall that c^{-1} denotes the conjugacy class $\{x^{-1}|x \in c\}$ of Γ, and ζ_c is the order of the centralizer of the class c in Γ. Sometimes we will also view the bilinear form as a function of $(t_1, t_2) \in \mathbb{C}^\times \times \mathbb{C}^\times$:

$$\langle f, g\rangle_\Gamma^{r,\,s}(t_1, t_2) = \sum_{c\in\Gamma_*} \zeta_c^{-1} f(c, (t_1, t_2)) S(g(c, (t_1, t_2))).$$

The following is a direct consequence of the orthogonality of irreducible characters of Γ.

$$\begin{aligned} \langle \gamma_i \otimes r^m s^n, \gamma_j \otimes r^k s^l\rangle_\Gamma^{r,\,s} &= \delta_{ij} r^{m-k} s^{n-l}, \\ \sum_{\gamma\in\Gamma^*} \gamma(c') S(\gamma)(c) &= \delta_{c,c'}\zeta_c, \quad c, c' \in \Gamma_*. \end{aligned} \tag{3.1}$$

Let $\langle\ ,\ \rangle_{\Gamma_n}^{r,\,s}$ be the $\mathbb{C}[r^{\pm1}, s^{\pm1}]$-valued bilinear form on $R(\Gamma_n \times \mathbb{C}^\times \times \mathbb{C}^\times)$. The $\mathbb{C}[r^{\pm1}, s^{\pm1}]$-valued standard bilinear form in $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ is defined in terms of the bilinear form on $R(\Gamma_n \times \mathbb{C}^\times \times \mathbb{C}^\times)$ as follows:

$$\langle u, v\rangle^{r,\,s} = \sum_{n\geq 0} \langle u_n, v_n\rangle_{\Gamma_n}^{r,\,s},$$

where $u = \sum_n u_n$ and $v = \sum_n v_n$ with $u_n, v_n \in R(\Gamma_n \times \mathbb{C}^\times \times \mathbb{C}^\times)$.

3.2. A weighted bilinear form on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$. A class function $\xi \in R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ is called self-dual if for all $x \in \Gamma, (t_1, t_2) \in \mathbb{C}^\times \times \mathbb{C}^\times$

$$\xi(x, (t_1, t_2)) = S(\xi(x, (t_1, t_2))),$$

or equivalently $\xi^{r,\,s}(x) = \xi^{r^{-1},\,s^{-1}}(x^{-1})$.

We fix a self-dual class function ξ. The tensor product of two representations γ and β in $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times})$ will be denoted by $\gamma * \beta$.

Let $a_{ij} \in \mathbb{C}[r^{\pm1}, s^{\pm1}]$ be the (virtual) multiplicity of γ_j in $\xi * \gamma_i$, i.e.,

$$\xi * \gamma_i = \sum_{j=0}^{r} a_{ij}\gamma_j. \tag{3.2}$$

We denote by $A^{r,\,s}$ the $n \times n$ matrix $(a_{ij})_{0\leq i,j\leq n-1}$.

Associated to ξ we introduce the following weighted bilinear form

$$\langle f, g\rangle_\xi^{r,\,s} = \langle \xi * f, g\rangle_\Gamma^{r,\,s}, \quad f, g \in R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}.$$

where we use the superscript $r\,s$ to indicate the r, s-dependence. The superscript $r\,s$ is often omitted if the r, s-variable in characters f and g is clear from the context. The explicit formula of the bilinear form is given as follows.

$$\begin{aligned} \langle f, g\rangle_\xi^{r,\,s} &= \frac{1}{|\Gamma|}\sum_{x\in\Gamma} \xi^{r,\,s}(x) f^{r,\,s}(x) g^{r^{-1},\,s^{-1}}(x^{-1}) \\ &= \sum_{c\in\Gamma_*} \zeta_c^{-1} \xi^{r,\,s}(c) f^{r,\,s}(c) g^{r^{-1},\,s^{-1}}(c^{-1}), \end{aligned} \tag{3.3}$$

which is the average of the character $\xi * f * \overline{g}$ over Γ.

The self-duality of ξ together with (3.3) implies that

$$a_{ij} = \overline{a_{ji}},$$

i.e. $A^{r\,s}$ is a hermitian-like matrix with the bar action given by $\overline{r} = s, \overline{s} = r$.

The orthogonality (3.1) implies that

$$a_{ij} = \langle \gamma_i, \gamma_j \rangle_{\xi}^{r,\, s}. \tag{3.4}$$

Remark 3.1. If ξ is the trivial character γ_0, then the weighted bilinear form becomes the standard one on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

3.3. A weighted bilinear form on $R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)$. Let V be a $\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times$-module which affords a character γ in $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$. We can decompose V as follows:

$$V = \bigoplus_i V_i \otimes \mathbb{C}(k_i, l_i),$$

where V_i is a (virtual) Γ-module in $R(\Gamma)$ and $\mathbb{C}(k_i, l_i)$ is the one dimensional $\mathbb{C}^\times\times\mathbb{C}^\times$-module afforded by the character $r^{k_i}s^{l_i}$.

The n-th outer tensor product $V^{\otimes n}$ of V can be regarded naturally as a representation of the wreath product $(\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times)_n$ via permutation of the factors and the usual direct product action. More precisely, note that $\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times$ can be viewed as a subgroup of $(\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times)_n$ by the diagonal inclusion from $\mathbb{C}^\times\times\mathbb{C}^\times$ to $(\mathbb{C}^\times\times\mathbb{C}^\times)^n$:

$$\Gamma_n \times \mathbb{C}^\times\times\mathbb{C}^\times \longrightarrow (\Gamma^n \times {\mathbb{C}^\times\times\mathbb{C}^\times}^n) \rtimes S_n = (\Gamma \times \mathbb{C}^\times\times\mathbb{C}^\times)_n.$$

This provides a natural $\Gamma_n \times \mathbb{C}^\times\times\mathbb{C}^\times$-module structure on $V^{\otimes n}$. We denote its character by $\eta_n(\gamma)$. Explicitly we have

$$(g, \sigma, (t_1, t_2)).(v_1 \otimes \cdots \otimes v_n) = (g_1, (t_1, t_2))v_{\sigma^{-1}(1)} \otimes \cdots \otimes (g_n, (t_1, t_2))v_{\sigma^{-1}(n)}, \tag{3.5}$$

where $g = (g_1, \ldots, g_n) \in \Gamma^n$.

Let ε_n be the (1-dimensional) sign representation of Γ_n so that Γ^n acts trivially while letting S_n act as a sign representation. We denote by $\varepsilon_n(\gamma) \in R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)$ the character of the tensor product of $\varepsilon_n \otimes 1$ and $V^{\otimes n}$.

The weighted bilinear form on $R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)$ is now defined by

$$\langle f, g\rangle_{\xi,\Gamma_n}^{r,\, s} = \langle \eta_n(\xi) * f, g\rangle_{\Gamma_n}^{r,\, s}, \quad f, g \in R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times).$$

We shall see in Corollary 6.4 that $\eta_n(\xi)$ is self-dual if the class function ξ is invariant under the antipode S. In such a case the matrix of the bilinear form $\langle\ ,\ \rangle_{\xi}^{r,\, s}$ is equal to its adjoint (transpose and bar action).

We can naturally extend η_n to a map from $R(\Gamma) \otimes r^m s^n$ to $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ as follows. In particular, if β and γ are characters of representations V and W of Γ respectively, then

$$\begin{aligned}&\eta_n(\beta \otimes r^m s^n + \gamma \otimes r^k s^l)\\ &= \sum_{m=0}^{n} Ind_{\Gamma_{n-m}\times\mathbb{C}^\times\times\mathbb{C}^\times\times\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times}^{\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times}[\eta_{n-m}(\beta \otimes r^m s^n) \otimes \eta_m(\gamma \otimes r^k s^l)],\end{aligned} \tag{3.6}$$

$$\begin{aligned}&\eta_n(\beta \otimes r^m s^n - \gamma \otimes r^k s^l)\\ &= \sum_{m=0}^{n} (-1)^m Ind_{\Gamma_{n-m}\times\mathbb{C}^\times\times\mathbb{C}^\times\times\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times}^{\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times}[\eta_{n-m}(\beta \otimes r^m s^n) \otimes \varepsilon_m(\gamma \otimes r^k s^l)].\end{aligned} \tag{3.7}$$

On $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times} = \bigoplus_n R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)$ the weighted bilinear form is given by

$$\langle u, v\rangle_{\xi}^{r,\, s} = \sum_{n\geq 0} \langle u_n, v_n\rangle_{\xi,\Gamma_n}^{r\, s}$$

where $u = \sum_n u_n$ and $v = \sum_n v_n$ with $u_n, v_n \in R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)$.

The bilinear form $\langle\ ,\ \rangle_\xi^{r,\,s}$ on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ is $\mathbb{C}$-bilinear and takes values in $\mathbb{C}[r^{\pm 1}, s^{\pm 1}]$. When $n = 1$, it reduces to the weighted bilinear form defined on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

We will often omit the superscript $r\,s$ and use the notation $\langle\ ,\ \rangle_\xi$ for the weighted bilinear form on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

4. Two-parameter quantum McKay weights

4.1. Two-parameter quantum McKay correspondence. Let $d_i = \gamma_i(c^0)$ be the dimension of the irreducible representation of Γ corresponding to the character γ_i.

The following generalizes a result of McKay [**Mc**].

Proposition 4.1. *For each class $c \in \Gamma_*$ the column vector*

$$v(c) = (\gamma_0(c), \gamma_1(c), \dots, \gamma_{n-1}(c))^t$$

is an eigenvector of the $n\times n$-matrix $A^{r\,s} = (\langle\gamma_i, \gamma_j\rangle_\xi^{r,\,s})$ with eigenvalue $\xi^{r,\,s}(c)$. In particular $(d_0, d_1, \dots, d_{n-1})$ is an eigenvector of $A^{r,\,s}$ with eigenvalue $\xi^{r,\,s}(c^0)$.

Proof. We compute directly that

$$\begin{aligned}\sum_{k=0}^{r}\langle\gamma_i, \gamma_k\rangle_\xi^{r,\,s}\gamma_k(c) &= \sum_k\sum_{c'\in\Gamma_*}\zeta_{c'}^{-1}\xi^{r,\,s}(c')\gamma_i(c')\gamma_k({c'}^{-1})\gamma_k(c)\\ &= \sum_{c'\in\Gamma_*}\zeta_{c'}^{-1}\xi^{r,\,s}(c')\gamma_i(c')\sum_k\gamma_k({c'}^{-1})\gamma_k(c)\\ &= \sum_{c'\in\Gamma_*}\zeta_{c'}^{-1}\xi^{r,\,s}(c')\gamma_i(c')\zeta_c\delta_{cc'}\\ &= \xi^{r,\,s}(c)\gamma_i(c).\end{aligned}$$

□

Let π be an irreducible faithful representation π of Γ of dimension d. For each integer n we define the r, s-integer $[n]$ that can be viewed as a character of $\mathbb{C}^\times\times\mathbb{C}^\times$ by

$$[n] = \frac{r^n - s^n}{r - s} = r^{n-1} + r^{n-2}s + \cdots + rs^{n-2} + s^{n-1}. \tag{4.1}$$

We take the following special class function

$$\xi = \gamma_0 \otimes [d](rs)^{-\frac{d}{4}} - \pi \otimes 1_{\mathbb{C}^\times\times\mathbb{C}^\times}, \tag{4.2}$$

where we have also used the symbol π for the corresponding character, and $1_{\mathbb{C}^\times\times\mathbb{C}^\times} = r^0s^0$ is the trivial character of $\mathbb{C}^\times\times\mathbb{C}^\times$.

Similar to one-parameter quantum case, we have the following fact(cf. [**FJW2**]).

Proposition 4.2. *The weighted bilinear form associated to (4.2) is non-degenerate. If π is an embedding of Γ into SU_d and $t \neq 1$ is a nonnegative real number, then the weighted bilinear form evaluated on t is positive definite.*

Remark 4.3. The matrix $A^{1,\,1}$ is integral, and the entries of $A^{r,\,s}$ are the r, s-numbers of the corresponding entries in $A^{1,\,1}$ when $r \geq 2$.

4.2. Two quantum McKay weights. Let Γ is a finite subgroup of SU_2 and we introduce the first distinguished self-dual class function

$$\xi = \gamma_0 \otimes ((rs^{-1})^{\frac{1}{2}} + (r^{-1}s)^{\frac{1}{2}}) - \pi \otimes 1_{\mathbb{C}^\times \times \mathbb{C}^\times},$$

where π is the character of the embedding of Γ in SU_2.

The matrix of the weighted bilinear form $\langle\ ,\ \rangle_\xi$ (cf. (3.4)) has the following entries:

$$a_{ij} = \begin{cases} (rs^{-1})^{\frac{1}{2}} + (r^{-1}s)^{\frac{1}{2}}, & \text{if } i = j, \\ -1, & \text{if } \langle\gamma_i, \gamma_j\rangle_\xi^1 = -1, \end{cases} \tag{4.3}$$

In particular when $r = s = 1$ the matrix $(a_{ij}^{1,\,1})$ coincides with the extended Cartan matrix of ADE type according to the five classes of finite subgroups of SU_2: the cyclic, binary dihedral, tetrahedral, octahedral, and icosahedral groups. McKay [**Mc**] gave a direct correspondence between a finite subgroup of SU_2 and the affine Dynkin diagram D of ADE type. Each irreducible character γ_i corresponds to a vertex of D, and the number of edges between γ_i and γ_j $(i \neq j)$ is equal to $|\langle\gamma_i, \gamma_j\rangle_\xi^{1,\,1}|$, where $\langle\gamma_i, \gamma_j\rangle_\xi^{1,\,1} = a_{ij}^1$ are the entries of matrix $A^{1\,1}$ of the weighted bilinear form $\langle\ ,\ \rangle_\xi^{1,\,1}$. For this reason we will call our matrix $A^{r,\,s} = (a_{ij}) = (\langle\gamma_i, \gamma_j\rangle_\xi^{r,\,s})$ the quantum Cartan matrix.

5. Two-parameter quantum Heisenberg algebras and Γ_n

5.1. Two-parameter Heisenberg algebra $\widehat{\mathfrak{h}}_{\Gamma,\xi}$. Let $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ be the infinite dimensional Heisenberg algebra over $\mathbb{C}[r^{\pm 1}, s^{\pm 1}]$, associated with Γ and $\xi \in R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$, with generators $a_m(c), c \in \Gamma_*, m \in \mathbb{Z}$ and a central element C subject to the following commutation relations:

$$[a_m(c^{-1}), a_n(c')] = m\delta_{m,-n}\delta_{c,c'}\zeta_c\xi_{r^m,\,s^m}(c)C, \quad c, c' \in \Gamma_*. \tag{5.1}$$

For $m \in \mathbb{Z}, \gamma \in \Gamma^*$ and $k,\, l \in \mathbb{Z}$ we define

$$a_m(\gamma \otimes r^k s^l) = \sum_{c\in\Gamma_*} \zeta_c^{-1}\gamma(c)a_m(c)r^{mk}s^{ml}$$

and then extend it to $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ linearly over $\mathbb{C}$. Thus we have for $\gamma \in R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$

$$a_m(\gamma) = \sum_{c\in\Gamma_*} \zeta_c^{-1}\gamma_{r^m,\,s^m}(c)a_m(c). \tag{5.2}$$

In particular we have $a_m(\gamma \otimes r^k s^l) = a_m(\gamma)r^{mk}s^{ml}$.

It follows immediately from the orthogonality (3.1) of the irreducible characters of Γ that for each $c \in \Gamma_*$

$$a_m(c) = \sum_{\gamma\in\Gamma^*} S(\gamma(c))a_m(\gamma).$$

Note that this formula is also valid if the summation runs through $\Gamma^* \otimes r^k s^l$ with a fixed k and l.

Proposition 5.1. *The Heisenberg algebra $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ has a new basis given by $a_n(\gamma)$ and C $(n \in \mathbb{Z}, \gamma \in \Gamma^*)$ over $\mathbb{C}[r^{\pm 1}, s^{\pm 1}]$ with the following relations:*

$$[a_m(\gamma), a_n(\gamma')] = m\delta_{m,-n}\langle\gamma, \gamma'\rangle_\xi^{r^m,\,s^m}C. \tag{5.3}$$

Proof. This is proved by a direct computation using Eqns. (5.1), (3.3) and (3.1).

$$\begin{aligned}
[a_m(\gamma), a_n(\gamma')] &= \sum_{c,c'\in\Gamma_*} \zeta_c^{-1}\zeta_{c'}^{-1}\gamma(c)\gamma'(c')[a_m(c), a_n(c')]\\
&= m\delta_{m,-n}\sum_{c,c'\in\Gamma_*} \zeta_c^{-1}\zeta_{c'}^{-1}\gamma(c)\gamma'(c')\delta_{c^{-1},c'}\zeta_c\xi_{r^m,\,s^m}(c)C\\
&= m\delta_{m,-n}\sum_{c\in\Gamma_*} \zeta_c^{-1}\gamma(c)\gamma'(c^{-1})\xi_{r^m,\,s^m}(c)C\\
&= m\delta_{m,-n}\langle\gamma,\gamma'\rangle_\xi^{r^m,\,s^m}C.
\end{aligned}$$

□

5.2. Action of $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ on the Space $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$. Let $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ be the symmetric algebra generated by $a_{-n}(\gamma), n\in\mathbb{N}, \gamma\in\Gamma_*$ over $\mathbb{C}[r^{\pm1}, s^{\pm1}]$. We define $a_{-n}(\gamma\otimes r^ks^l) = a_{-n}(\gamma)r^{-kn}s^{-ln}$ and the natural degree operator on the space $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ by

$$\deg(a_{-n}(\gamma\otimes r^ks^l)) = n$$

which makes $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ into a $\mathbb{Z}_+$-graded algebra.

The space $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ affords a natural realization of the Heisenberg algebra $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ with $C = 1$. Since $a_{-n}(\gamma\otimes r^ks^l) = r^{-nk}s^{-nl}a_{-n}(\gamma)$, it is enough to describe the action for $a_{-n}(\gamma)$. The central element C acts as the identity operator. For $n > 0$, $a_{-n}(\gamma)$ act as multiplication operators on $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$. The element $a_n(\gamma), n\geq 0$ acts as a differential operator through contraction:

$$\begin{aligned}
&a_n(\gamma).a_{-n_1}(\alpha_1)a_{-n_2}(\alpha_2)\dots a_{-n_k}(\alpha_k)\\
&= \textstyle\sum_{i=1}^k\langle\gamma,\alpha_i\rangle_\xi^{r^n,\,s^n}a_{-n_1}(\alpha_1)a_{-n_2}(\alpha_2)\dots\check{a}_{-n_i}(\alpha_i)\dots a_{-n_k}(\alpha_k).
\end{aligned}$$

Here $n_i > 0, \alpha_i\in R(\Gamma)$ for $i = 1,\dots,k$, and $\check{a}_{-n_i}(\alpha_i)$ means the very term is deleted. In this case $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ is an irreducible representation of $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ with the unit 1 as the highest weight vector.

5.3. The bilinear form on $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$. As a $\widehat{\mathfrak{h}}_{\Gamma,\xi}$-module, the space $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ admits a bilinear form $\langle\ ,\ \rangle'_\xi$ over $\mathbb{C}[r^{\pm1}, s^{\pm1}]$ characterized by

$$\begin{aligned}
&\langle 1,1\rangle'_\xi = 1,\\
&\langle au, v\rangle'_\xi = \langle u, a^*v\rangle'_\xi, \qquad a\in\widehat{\mathfrak{h}}_{\Gamma,\xi},
\end{aligned}\tag{5.4}$$

with the adjoint map $*$ on $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ given by

$$a_n(\gamma\otimes r^ks^l)^* = a_{-n}(\gamma\otimes r^ks^l), \qquad n\in\mathbb{Z}.\tag{5.5}$$

Note that the adjoint map $*$ is a $\mathbb{C}$-linear anti-homomorphism of $\widehat{\mathfrak{h}}_{\Gamma,\xi}$, and $r^* = \overline{r}, s^* = \overline{s}$. We still use the same symbol $*$ to denote the hermitian-like dual, since it clearly generalizes the $*$-action on the deformed Cartan matrix (**??**).

For any partition $\lambda = (\lambda_1, \lambda_2, \dots)$ and $\gamma\in\Gamma^*$, we define

$$a_{-\lambda}(\gamma) = a_{-\lambda_1}(\gamma)a_{-\lambda_2}(\gamma)\dots.$$

For $\rho = (\rho(\gamma))_{\gamma\in\Gamma^*}\in\mathcal{P}(\Gamma^*)$, we define

$$a_{-\rho\otimes r^ks^l} = r^{-k\|\rho\|}s^{-l\|\rho\|}\prod_{\gamma\in\Gamma^*}a_{-\rho(\gamma)}(\gamma).$$

It is clear that for a fixed $k \in \mathbb{Z}$ the elements $a_{-\rho\otimes q^k}, \rho \in \mathcal{P}(\Gamma^*)$ form a basis of $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ over $\mathbb{C}[r^{\pm1}, s^{\pm1}]$.

Given a partition $\lambda = (\lambda_1, \lambda_2, \dots)$ and $c \in \Gamma_*$, we define

$$a_{-\lambda}(c\otimes r^k s^l) \quad = \quad r^{-k|\lambda|}s^{-l|\lambda|}a_{-\lambda_1}(c)a_{-\lambda_2}(c)\dots,$$

For any $\rho = (\rho(c))_{c\in\Gamma_*} \in \mathcal{P}(\Gamma_*)$ and $k \in \mathbb{Z}$, we define

$$a'_{-\rho\otimes r^k s^l} = r^{-k\|\rho\|}s^{-l\|\rho\|}\prod_{c\in\Gamma_*} a_{-\rho(c)}(c).$$

It follows from Proposition 5.1 that

$$\langle a'_{-\rho\otimes r^m s^n}, a'_{-\overline{\sigma}\otimes r^k s^l}\rangle'_\xi = \delta_{\rho,\sigma} r^{\|\rho\|(k-m)} s^{\|\rho\|(l-n)} Z_\rho \prod_{c\in\Gamma_*}\prod_{i\geq 1}\xi_{q^i}(c)^{m_i(\rho(c))},$$

where $\rho, \sigma \in \mathcal{P}(\Gamma_*)$. Note that $S(a'_{-\rho\otimes r^k s^l}) = a'_{-\overline{\rho}\otimes r^{-k}s^{-l}}$, where we recall that $\overline{\rho} \in \mathcal{P}(\Gamma_*)$ is the partition-valued function given by $c \mapsto \rho(c^{-1})$, $c \in \Gamma$.

6. The characteristic map as an isometry

6.1. The characteristic map ch. Let $\Psi : \Gamma_n \longrightarrow S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ be the map defined by $\Psi(x) = a'_{-\rho}$ if $x \in \Gamma_n$ is of type ρ.

We define a $\mathbb{C}$-linear map $ch : R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times} \longrightarrow S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ by letting

$$\begin{aligned} ch(f) &= \langle f, \Psi\rangle_{\Gamma_n} \\ &= \sum_{\rho\in\mathcal{P}(\Gamma_*)} Z_\rho^{-1} S(f(\rho)) a'_{-\rho}, \end{aligned} \tag{6.1}$$

where $f(\rho) \in \mathbb{C}[r^{\pm1}, s^{\pm1}]$ is the value of f at the elements of type ρ. The map ch is called the *characteristic map*. This generalizes the definition of the characteristic map in the classical setting (cf. [**M, FJW1, FJW2**]).

The space $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ can also be interpreted as follows. The element $a_{-n}(\gamma), n > 0, \gamma \in \Gamma^*$ is identified as the n-th power sum in a sequence of variables $y_g = (y_{i\gamma})_{i\geq 1}$. By the commutativity among $a_{-n}(\gamma)$ $(\gamma \in \Gamma^*, n > 0)$ and dimension counting it is clear that the space $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ is isomorphic with the space Λ_Γ of symmetric functions indexed by Γ^* tensored with $\mathbb{C}[r^{\pm1}, s^{\pm1}]$ (cf. [**M**]).

Denote by $c_n (c \in \Gamma_*)$ the conjugacy class in Γ_n of elements $(x, q) \in \Gamma_n$ such that q is an n-cycle and $x \in c$. Denote by $\sigma_n(c \otimes r^k s^l)$ the class function on $\Gamma_n \times \mathbb{C}^\times \times \mathbb{C}^\times$ which takes values $n\zeta_c t_1^{-nk} t_2^{-nl}$ (i.e. the order of the centralizer of an element in the class c_n times $t_1^{-nk}t_2^{-nl}$) on elements in the class $c_n \times t_1 t_2$ and 0 elsewhere. For $\rho = \{m_r(c)\}_{r\geq 1, c\in\Gamma_*} \in \mathcal{P}_n(\Gamma^*)$ and $k \in \mathbb{Z}$,

$$\sigma_{\rho\otimes r^k s^l} = r^{nl}s^{nk}\prod_{a\geq 1, c\in\Gamma_*}\sigma_a(c)^{m_a(c)}$$

is the class function on $\Gamma_n \times \mathbb{C}^\times \times \mathbb{C}^\times$ which takes value $Z_\rho t_1^{-nk}t_2^{-nl}$ on the conjugacy class of type $\rho \times t_1 t_2$ and 0 elsewhere. Given $\gamma \in \Gamma^*$ and $k, l \in \mathbb{Z}$, we denote by $\sigma_n(\gamma \otimes r^k s^l)$ the class function on $\Gamma_n \times \mathbb{C}^\times \times \mathbb{C}^\times$ which takes values $n\gamma(c) t_1^{-nk} t_2^{-nl}$ on elements in the class $c_n \times t_1 t_2 (c \in \Gamma_*)$ and 0 elsewhere.

Lemma 6.1. *The map ch sends* $\sigma_{\rho\otimes r^k s^l}$ *to* $a'_{-\rho\otimes r^k s^l}$. *In particular, it sends* $\sigma_n(\gamma \otimes r^k s^l)$ *to* $a_{-n}(\gamma \otimes r^k s^l)$ *in* $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

Proof. This is verified by the definition of ch (6.1) and the character values of σ_n defined above. □

Proposition 6.2. *Given $\gamma \in \Gamma^*$, the character value of $\eta_n(\gamma \otimes r^k s^l)$ on the conjugacy class c_ρ of type $\rho = (\rho(c))_{c\in\Gamma_*}$ is given by*

$$\eta_n(\gamma \otimes r^k s^l)(c_\rho) = \prod_{c\in\Gamma_*} \gamma(c)^{l(\rho(c))} r^{nk} s^{nl}. \tag{6.2}$$

In particular, we have $\eta_n(\gamma \otimes r^k s^l) = \eta_n(\gamma) r^{nk} s^{nl}$.

Proof. We first let (g, σ) be an element of Γ_n such that σ is a cycle of length n, say $\sigma = (12\cdots n)$. Let $\{e_i\}$ be a basis of V, and $\gamma \otimes r^k s^l$ is afforded by the action: $(h,t)e_j = \sum_i c_{ij}(h) t^k e_i$, where $h \in \Gamma$. We then have

$$\begin{aligned}
&(g,\sigma,t).(e_{j_1}\otimes e_{j_2}\otimes\cdots\otimes e_{j_n})\\
&= (g_1,t)e_{j_n}\otimes(g_2,t)e_{j_1}\otimes\cdots\otimes(g_n,t)e_{j_{n-1}}\\
&= \sum_{i_1,\dots,i_n} t^{kn} c_{i_nj_n}(g_1)c_{i_1j_1}(g_2)\cdots c_{i_{n-1}j_{n-1}}(g_n) e_{i_n}\otimes e_{i_1}\cdots\otimes e_{i_{n-1}}.
\end{aligned}$$

It follows that

$$\begin{aligned}
\eta_n(\gamma\otimes r^k s^l)(c_\rho,t) &= \text{trace } (g,\sigma,t)\\
&= \sum_{j_1,\dots,j_n} t^{kn} c_{j_1j_n}(g_1)c_{j_2j_1}(g_2)\cdots c_{j_nj_{n-1}}(g_n)\\
&= \text{trace } t^{kn} a(g_n)a(g_{n-1})\dots a(g_1)\\
&= \text{trace } t^{kn} a(g_ng_{n-1}\dots g_1) = \gamma(c) r^{kn} s^{ln}(t).
\end{aligned}$$

Given $x \times y \in \Gamma_n$ where $x \in \Gamma_r$ and $y \in \Gamma_{n-r}$, by (3.5) we clearly have

$$\eta_n(\gamma\otimes r^k s^l)(x\times y, t) = \eta_n(\gamma\otimes r^k s^l)(x,t)\eta_n(\gamma\otimes r^k s^l)(y,t).$$

This immediately implies the formula. □

A similar argument gives that

$$\varepsilon_n(\gamma\otimes r^k s^l)(x,t) = (-1)^n \prod_{c\in\Gamma_*}(-\gamma(c))^{l(\rho(c))} t^{nk}, \tag{6.3}$$

where x is any element in the conjugacy class of type $\rho = (\rho(c))_{c\in\Gamma^*}$.

Formula (6.2) is equivalent to the following:

$$\eta_n(\gamma\otimes r^k s^l)(c_\rho,t) = \prod_{c\in\Gamma_*}\prod_{i\geq 1}(\gamma\otimes r^k s^l)(c,t^i)^{m_i(\rho(c))}. \tag{6.4}$$

The following result allows us to extend the map from $\gamma \in \Gamma^*$ to $R(\Gamma_n)$.

Proposition 6.3. *For any $\gamma \in R(\Gamma)$, we have*

$$\sum_{n\geq 0} ch(\eta_n(\gamma\otimes r^k s^l)) z^n = \exp\left(\sum_{n\geq 1}\frac{1}{n} a_{-n}(\gamma)(r^{-k}s^{-l}z)^n\right), \tag{6.5}$$

$$\sum_{n\geq 0} ch(\varepsilon_n(\gamma\otimes r^k s^l)) z^n = \exp\left(\sum_{n\geq 1}(-1)^{n-1}\frac{1}{n} a_{-n}(\gamma)(r^{-k}s^{-l}z)^n\right). \tag{6.6}$$

PROOF. It follows from definition of ch (6.1) and (6.4) that

$$\begin{aligned}
&\sum_{n\geq 0} \mathrm{ch}(\eta_n(\gamma\otimes r^k s^l))z^n \\
=\ &\sum_{\rho} Z_{\rho}^{-1} \prod_{c\in\Gamma_*}\prod_{i\geq 1} S(\gamma_{r^{ik}s^{il}}(c)^{m_i(\rho(c))})a_{-\rho(c)} z^{||\rho||} r^{-k||\rho||} s^{-l||\rho||} \\
=\ &\sum_{\rho} Z_{\rho}^{-1} \prod_{c\in\Gamma_*} \gamma(c)^{l(\rho(c))} a_{-\rho(c)} (r^{-k}s^{-l}z)^{||\rho||} \\
=\ &\prod_{c\in\Gamma_*} \Big(\sum_{\lambda} (\zeta_c^{-1}\gamma(c))^{l(\lambda)} z_{\lambda}^{-1} a_{-\lambda}(c)(r^{-k}s^{-l}z)^{|\lambda|}\Big) \\
=\ &\exp\left(\sum_{n\geq 1}\frac{1}{n}\sum_{c\in\Gamma_*}\zeta_c^{-1}\gamma(c)a_{-n}(c)(r^{-k}s^{-l}z)^n\right) \\
=\ &\exp\left(\sum_{n\geq 1}\frac{1}{n}\, a_{-n}(\gamma)(r^{-k}s^{-l}z)^n\right).
\end{aligned}$$

Similarly we can prove (6.6) using the following identity

$$\begin{aligned}
\varepsilon_n(\gamma\otimes r^k s^l)(x) &= (-1)^n \prod_{c\in\Gamma_*}\prod_{i\geq 1} (-\gamma_{r^{ik}s^{il}}(c))^{m_i(\rho(c))} \\
&= (-r^k s^l)^n \prod_{c\in\Gamma_*}\prod_{i\geq 1}(-\gamma(c))^{m_i(\rho(c))} \\
&= \varepsilon_n(\gamma)(x) r^{nk}s^{nl}.
\end{aligned}$$

The same argument as in the classical case (cf. [**FJW1, FJW2**]) by using (3.6) and (3.7) will show that the proposition holds for linear combination of simple characters such as $\gamma\otimes r^k s^l - \beta\otimes q^k$, and thus it is true for any element $\gamma\otimes r^k s^l$, where $\gamma\in R(\Gamma)$. □

Comparing components we obtain

$$\begin{aligned}
\mathrm{ch}(\eta_n(\gamma\otimes r^k s^l)) &= \sum_{\lambda} \frac{r^{-kn}s^{-ln}}{z_{\lambda}}\, a_{-\lambda}(\gamma), \\
\mathrm{ch}(\varepsilon_n(\gamma\otimes r^k s^l)) &= \sum_{\lambda} \frac{r^{-kn}s^{-ln}}{z_{\lambda}}\, (-1)^{|\lambda|-l(\lambda)} a_{-\lambda}(\gamma),
\end{aligned}$$

where the sum runs over all partitions λ of n.

COROLLARY 6.4. The formula (6.4) remains valid when $\gamma\otimes r^k s^l$ is replaced by any element $\xi\in R(\Gamma\times\mathbb{C}^{\times})$. In particular $\eta_n(\xi)$ is self-dual provided that ξ is invariant under the antipode S.

6.2. Isometry between $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ and $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$. The symmetric algebra $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times} = S_{\Gamma}\otimes\mathbb{C}[r^{\pm 1}, s^{\pm 1}]$ has the following Hopf algebra structure over $\mathbb{C}$. The multiplication is the usual one, and the comultiplication is given by

$$\Delta(r^k s^l) = r^k s^l \otimes r^k s^l$$

$$\Delta(a_n(\gamma\otimes r^k s^l)) = a_n(\gamma\otimes r^k s^l)\otimes r^{nk}s^{nl} + r^{nk}s^{nl}\otimes a_n(\gamma\otimes r^k s^l),$$

where $\gamma \in \Gamma^*$. The last formula is equivalent to the following:

$$\Delta(a_n(c \otimes r^k s^l)) = a_n(c \otimes r^k s^l) \otimes r^{nk} s^{nl} + r^{nk} s^{nl} \otimes a_n(c \otimes r^k s^l), \tag{6.7}$$

where $c \in \Gamma_*$. The antipode is given by

$$S(r^k s^l) = r^l s^k,$$
$$S(a_n(\gamma \otimes r^k s^l)) = -a_n(\gamma \otimes r^l s^k)$$

The antipode commutes with the adjoint (dual) map $*$:

$$*^2 = S^2 = Id, \qquad S* = *S. \tag{6.8}$$

Recall that we have defined a Hopf algebra structure on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ in Sect. 2.

PROPOSITION 6.5. *The characteristic map* $ch : R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times} \longrightarrow S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ *is an isomorphism of Hopf algebras.*

PROOF. It follows immediately from the definition of the comultiplication in the both Hopf algebras (cf. (2.4) and (6.7)). □

Recall that we have defined a bilinear form $\langle\ ,\ \rangle_\xi$ on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ and a bilinear form on $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ denoted by $\langle\ ,\ \rangle'_\xi$, where ξ is a self-dual class function. The following lemma is immediate from our definition of $\langle\ ,\ \rangle'_\xi$ and the comultiplication Δ.

LEMMA 6.6. *The bilinear form* $\langle\ ,\ \rangle'_\xi$ *on* $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ *can be characterized by the following two properties:*

1). $\langle a_{-n}(\beta\otimes r^{k_1}s^{l_1}), a_{-m}(\gamma\otimes r^{k_2}s^{l_2})\rangle'_\xi = \delta_{n,m} r^{n(k_2-k_1)} s^{n(l_2-l_1)} \langle\beta,\gamma\rangle'_\xi$, $\quad \beta,\gamma \in \Gamma^*$, $k_1, k_2, l_1, l_2 \in \mathbb{Z}$.

2). $\langle fg, h\rangle'_\xi = \langle f \otimes g, \Delta h\rangle'_\xi$, *where* $f, g, h \in S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$, *and the bilinear form on* $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times} \otimes S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$, *is induced from* $\langle\ ,\ \rangle'_\xi$ *on* $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

THEOREM 6.7. *The characteristic map is an isometry from the space* $(R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}, \langle\ ,\ \rangle_\xi)$ *to the space* $(S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}, \langle\ ,\ \rangle'_\xi)$.

PROOF. By Corollary 6.4, the character value of $\eta_n(\xi)$ at an element x of type ρ is

$$\eta_n(\xi)(x) = \prod_{c\in\Gamma_*}\prod_{i\geq 1} \xi_{q^i}(c)^{m_i(\rho(c))}.$$

Thus it follows from definition that

$$\begin{aligned}
\langle \sigma_{\rho\otimes r^{k_1}s^{l_1}}, \sigma_{\rho'\otimes r^{k_2}s^{l_2}}\rangle_\xi &= \sum_{\mu\in\mathcal{P}_n(\Gamma_*)} Z_\mu^{-1} r^{n(k_2-k_1)} s^{n(l_2-l_1)} \xi_q(c_\mu)\sigma_\rho(c_\mu)\sigma_{\rho'}(c_\mu)\\
&= \delta_{\rho,\rho'} Z_\rho^{-1} r^{n(k_2-k_1)} s^{n(l_2-l_1)} \xi(c_\rho) Z_\rho Z_\rho\\
&= \delta_{\rho,\rho'} Z_\rho r^{n(k_2-k_1)} s^{n(l_2-l_1)} \prod_{c\in\Gamma_*}\prod_{i\geq 1} \xi_{q^i}(c)^{m_i(\rho(c))}.
\end{aligned}$$

By Lemma 6.1 and the formula (5.6), we see that

$$\langle \sigma_{\rho\otimes r^{k_1}s^{l_1}}, \sigma_{\rho'\otimes r^{k_2}s^{l_2}}\rangle_\xi = \langle a_{-\rho\otimes r^{k_1}s^{l_1}}, a_{-\rho'\otimes r^{k_2}s^{l_2}}\rangle'_\xi = \langle \mathrm{ch}(\sigma_{\rho\otimes r^{k_1}s^{l_1}}), \mathrm{ch}(\sigma_{\rho'\otimes r^{k_2}s^{l_2}})\rangle'_\xi.$$

Since $\sigma_{\rho\otimes r^k s^l}, \rho \in \mathcal{P}(\Gamma_*)$ form a $\mathbb{C}$-basis of $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$, we have shown that ch : $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times} \longrightarrow S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ is an isometry. □

From now on we will not distinguish the bilinear form $\langle\ ,\ \rangle_\xi$ on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ from the bilinear form $\langle\ ,\ \rangle'_\xi$ on $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

7. Two-parameter quantum vertex operators and $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$

7.1. Two-parameter vertex operators and Heisenberg algebras in $\mathcal{F}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

Let $\mathbb{K}=\mathbb{Q}(r,s)$ denote a field of rational functions with two-parameters r, s $(r\neq\pm s)$. Let Q be an integral lattice with basis α_i, $i=0,1,\ldots,n-1$ endowed with a symmetric bilinear form. We fix a 2-cocycle $\epsilon: Q\times Q\longrightarrow\mathbb{K}^\times$ has the following properties:

$$\epsilon(\alpha+\beta,\,\gamma)=\epsilon(\alpha,\,\beta)\epsilon(\beta,\,\gamma),$$
$$\epsilon(\alpha,\,\beta+\gamma)=\epsilon(\alpha,\,\beta)\epsilon(\alpha,\,\gamma)$$

We construct such a cocycle directly by

$$\epsilon(\alpha_i,\alpha_j)=\begin{cases}(-r_is_i)^{\frac{a_{ij}}{2}}, & i>j;\\ (rs)^{\frac{1}{2}}, & i=j;\\ 1, & i<j.\end{cases}$$

Let ξ be a self-dual virtual character in $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$. Recall that the lattice $R_{\mathbb{Z}}(\Gamma)$ is a $\mathbb{Z}$-lattice under the bilinear form $\langle\ ,\ \rangle_\xi^1$, here the superscript means $r=s^{-1}=1$. For our purpose we will always associate a 2-cocycle ϵ as in the previous subsection to the integral lattice $(R_{\mathbb{Z}}(\Gamma),\langle\ ,\ \rangle_\xi^1)$ (and its sublattices).

Let $\mathbb{C}[R_{\mathbb{Z}}(\Gamma)]$ be the group algebra generated by e^γ, $\gamma\in R_{\mathbb{Z}}(\Gamma)$. We introduce two special operators acting on $\mathbb{C}[R_{\mathbb{Z}}(\Gamma)]$: A ($\epsilon$-twisted) multiplication operator e^α defined by

$$e^\alpha.e^\beta=\epsilon(\alpha,\beta)e^{\alpha+\beta},\quad \alpha,\beta\in R_{\mathbb{Z}}(\Gamma),$$

and a differentiation operator ∂_α given by

$$\partial_\alpha e^\beta=\langle\alpha,\beta\rangle_\xi^1 e^\beta,\quad \alpha,\beta\in R_{\mathbb{Z}}(\Gamma).$$

These two operators are then extended linearly to the space

$$\mathcal{F}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}=R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}\otimes\mathbb{C}[R_{\mathbb{Z}}(\Gamma)]\tag{7.1}$$

by letting them act on the $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ part trivially.

We define the Hopf algebra structure on $\mathbb{C}[R_{\mathbb{Z}}(\Gamma)]$ and extend the Hopf algebra structure from $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ to $\mathcal{F}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ as follows.

$$\Delta(e^\alpha)=e^\alpha\otimes e^\alpha,\qquad S(e^\alpha)=e^{-\alpha}.$$

The bilinear form $\langle\ ,\ \rangle_\xi^{r\,s}$ on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ is extended to $\mathcal{F}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ by

$$\langle e^\alpha,e^\beta\rangle_\xi=\delta_{\alpha,\beta}.$$

With respect to this extended bilinear form we have the $*$-action (adjoint action) on the operators e^α and ∂_α:

$$(e^\alpha)^*=e^{-\alpha},\qquad (z^{\partial_\alpha})^*=z^{-\partial_\alpha}.\tag{7.2}$$

For each $k\in\mathbb{Z}$, we introduce the group theoretic operators $H_{\pm n}(\gamma\otimes r^ks^l), E_{\pm n}(\gamma\otimes r^ks^l), \gamma\in R(\Gamma), n>0$ as the following compositions of maps:

$$\begin{aligned}
H_{-n}(\gamma\otimes r^ks^l) \quad &: \quad R(\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times) \overset{\eta_n(\gamma\otimes r^ks^l)\otimes}{\longrightarrow} R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)\otimes R(\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times)\\
&\quad \overset{Ind\otimes m_{\mathbb{C}^\times}}{\longrightarrow} R(\Gamma_{n+m}\times\mathbb{C}^\times\times\mathbb{C}^\times)\\
E_{-n}(\gamma\otimes r^ks^l) \quad &: \quad R(\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times) \overset{\varepsilon_n(\gamma\otimes r^ks^l)\otimes}{\longrightarrow} R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)\otimes R(\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times)\\
&\quad \overset{Ind\otimes m_{\mathbb{C}^\times}}{\longrightarrow} R(\Gamma_{n+m}\times\mathbb{C}^\times\times\mathbb{C}^\times)\\
E_{n}(\gamma\otimes r^ks^l) \quad &: \quad R(\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times) \xrightarrow{Res} R(\Gamma_n)\otimes R(\Gamma_{m-n}\times\mathbb{C}^\times\times\mathbb{C}^\times)\\
&\quad \overset{\langle\varepsilon_n(\gamma\otimes r^ks^l),\cdot\rangle_\xi}{\longrightarrow} R(\Gamma_{m-n}\times\mathbb{C}^\times\times\mathbb{C}^\times)\\
H_{n}(\gamma\otimes r^ks^l) \quad &: \quad R(\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times) \xrightarrow{Res} R(\Gamma_n)\otimes R(\Gamma_{m-n}\times\mathbb{C}^\times\times\mathbb{C}^\times)\\
&\quad \overset{\langle\eta_n(\gamma\otimes q^k),\cdot\rangle_\xi}{\longrightarrow} R(\Gamma_{m-n}\times\mathbb{C}^\times\times\mathbb{C}^\times),
\end{aligned}$$

where Res and Ind are the restriction and induction functors in $R_\Gamma=\bigoplus_{n\geq 0}R(\Gamma_n)$.

We introduce their generating functions in a formal variable z:

$$\begin{aligned}
H_\pm(\gamma\otimes r^ks^l,z) &= \sum_{n\geq 0} H_{\mp n}(\gamma\otimes r^ks^l)z^{\pm n},\\
E_\pm(\gamma\otimes r^ks^l,z) &= \sum_{n\geq 0} E_{\mp n}(\gamma\otimes r^ks^l)(-z)^{\pm n}.
\end{aligned}$$

We now define the vertex operators $Y_n^\pm(\gamma\otimes r^ks^l,a,b)$, $\gamma\in\Gamma^*$, $k,l,a,b\in\mathbb{Z}$, $n\in\mathbb{Z}+\langle\gamma,\gamma\rangle_\xi^1/2$ as follows.

$$\begin{aligned}
Y^+(\gamma\otimes r^ks^l,a,b,z) &= \sum_{n\in\mathbb{Z}+\langle\gamma,\gamma\rangle_\xi^1/2} Y_n^+(\gamma\otimes r^ks^l,a,b)z^{-n-\langle\gamma,\gamma\rangle_\xi^1/2}\\
&= H_+(\gamma\otimes r^ks^l,z)E_-(\gamma\otimes r^{k-a}s^{l-b},z)e^\gamma(r^{-k}s^{-l}z)^{\partial_\gamma}, \qquad (7.3)
\end{aligned}$$

$$\begin{aligned}
Y^-(\gamma\otimes r^ks^l,a,b,z) &= (Y^+(\gamma\otimes r^ks^l,a,b,z^{-1}))^*\\
&= \sum_{n\in\mathbb{Z}+\langle\gamma,\gamma\rangle_\xi^1/2} Y_n^-(\gamma\otimes r^ks^l,a,b)z^{-n-\langle\gamma,\gamma\rangle_\xi^1/2}\\
&= E_+(\gamma\otimes r^{k-a}s^{l-b},z)H_-(\gamma\otimes r^ks^l,z)e^{-\gamma}(r^{-k}s^{-l}z)^{-\partial_\gamma}. \qquad (7.4)
\end{aligned}$$

One easily sees that the operators $Y_n^\pm(\gamma\otimes r^ks^l,a,b)$ are well-defined operators acting on the space $\mathcal{F}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

We extend the $\mathbb{Z}_+$-gradation on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ to a $\frac{1}{2}\langle\gamma,\gamma\rangle_\xi^1+\mathbb{Z}_+$-gradation on $\mathcal{F}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ by letting

$$\deg a_{-n}(\gamma\otimes r^ks^l)=n,\quad \deg e^\gamma=\frac{1}{2}\langle\gamma,\gamma\rangle_\xi^1.$$

We denote by $\overline{R}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ the subalgebra of $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ excluding the generators $a_n(\gamma_0)$, $n\in\mathbb{Z}^\times$. The bilinear form $\langle\ ,\ \rangle_\xi$ on

$$\overline{\mathcal{F}}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}=\overline{R}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}\otimes\overline{R}_{\mathbb{Z}}(\Gamma)$$

will be the restriction of $\langle\ ,\ \rangle_\xi$ on $\mathcal{F}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ to $\overline{\mathcal{F}}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

We define $\widetilde{a}_{-n}(\gamma\otimes r^k s^l), n>0$ to be a map from $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ to itself by the following composition

$$\begin{aligned} R(\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times) &\overset{\sigma_n(\gamma\otimes r^k s^l)\otimes}{\longrightarrow} R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)\otimes R(\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times)\\ &\overset{Ind\otimes m_{\mathbb{C}^\times}}{\longrightarrow} R(\Gamma_{n+m}\times\mathbb{C}^\times\times\mathbb{C}^\times). \end{aligned}$$

We also define $\widetilde{a}_n(\gamma\otimes r^k s^l), n>0$ to be a map from $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ to itself as the composition

$$\begin{aligned} R(\Gamma_m\times\mathbb{C}^\times\times\mathbb{C}^\times) &\overset{Res\otimes 1}{\longrightarrow} R(\Gamma_n\times\mathbb{C}^\times\times\mathbb{C}^\times)\otimes R(\Gamma_{m-n}\times\mathbb{C}^\times\times\mathbb{C}^\times)\\ &\overset{\langle\sigma_n(\gamma\otimes r^k s^l),\cdot\rangle_\xi^q}{\longrightarrow} R(\Gamma_{m-n}\times\mathbb{C}^\times\times\mathbb{C}^\times). \end{aligned}$$

Proposition 7.1. *The operators $\widetilde{a}_n(\gamma)$, $\gamma\in\Gamma^*, n\in\mathbb{Z}^\times$ satisfy the Heisenberg algebra relations (5.1) with $C=1$.*

Proof. This is similarly proved as for the classical setting in [**W**]. □

7.2. Group theoretic interpretation of vertex operators. To compare the vertex operators $Y^{\pm}(\gamma\otimes q^k,a,b,z)$ with the familiar vertex operators acting in the Fock space we introduce the space

$$V_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}=S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}\otimes\mathbb{C}[R_{\mathbb{Z}}(\Gamma)].$$

We extend the bilinear form $\langle\ ,\ \rangle_\xi^{r,s}$ in $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ to the space $V_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ and also extend the $\mathbb{Z}_+$-gradation on $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ to a $\frac{1}{2}\mathbb{Z}_+$-gradation on V_Γ.

We extend the characteristic map to the map

$$ch:\mathcal{F}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}\longrightarrow V_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$$

by identity on $R_{\mathbb{Z}}(\Gamma)$. Then Proposition 6.5 and Theorem 6.7 imply that we have an isometric isomorphism of Hopf algebras. We can now identify the operators from the previous subsections with the operators constructed from the Heisenberg algebra.

Theorem 7.2. *For any $\gamma\in R(\Gamma)$ and $k\in\mathbb{Z}$, we have*

$$ch\big(H_+(\gamma\otimes r^k s^l,z)\big) = \exp\Bigg(\sum_{n\geq 1}\frac{1}{n}\,a_{-n}(\gamma)(r^{-k}s^{-l}z)^n\Bigg), \tag{7.5}$$

$$ch\big(E_+(\gamma\otimes r^k s^l,z)\big) = \exp\Bigg(-\sum_{n\geq 1}\frac{1}{n}\,a_{-n}(\gamma)(r^{-k}s^{-l}z)^n\Bigg), \tag{7.6}$$

$$ch\big(H_-(\gamma\otimes r^k s^l,z)\big) = \exp\Bigg(\sum_{n\geq 1}\frac{1}{n}\,a_{n}(\gamma)(r^{-k}s^{-l}z)^{-n}\Bigg), \tag{7.7}$$

$$ch\big(E_-(\gamma\otimes r^k s^l,z)\big) = \exp\Bigg(-\sum_{n\geq 1}\frac{1}{n}\,a_{n}(\gamma)(r^{-k}s^{-l}z)^{-n}\Bigg). \tag{7.8}$$

Proof. The first and second identities were essentially established in Proposition 6.3 together with Lemma 6.1, where the components are viewed as operators acting on $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ or $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$. Note that $a_n(\gamma\otimes r^k s^l)=a_n(\gamma)r^{kn}s^{ln}$.

We observe from definition that the adjoint $*$-action of $E_+(\gamma\otimes r^k s^l,z)$ and $H_-(\gamma\otimes r^k s^l,z)$ with respect to the bilinear form $\langle\ ,\ \rangle_\xi^{r,s}$ are $E_-(\gamma\otimes r^k s^l,z^{-1})$

and $H_-(\gamma\otimes r^k s^l, z^{-1})$ respectively. The third and fourth identities are obtained by applying the adjoint action $*$ to the first two identities. □

Remark 7.3. Replacing γ by $-\gamma$ in (7.5) and (7.7) we obtain the equivalent formulas (7.6) and (7.8) respectively.

Applying the characteristic map to the vertex operators $Y^{\pm}(\gamma, a, b, z)$, we obtain the following group theoretical explanation of vertex operators acting on the Fock space $\mathcal{F}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

Theorem 7.4. *For any $\gamma \in R_\Gamma$ and $k \in \mathbb{Z}$, we have*

$$\begin{aligned}
&Y^+(\gamma, a, b, z)\\
&= \exp\Big(\sum_{n\geq 1}\frac{1}{n}\,\widetilde{a}_{-n}(\gamma)z^n\Big)\ \exp\Big(-\sum_{n\geq 1}\frac{1}{n}\,\widetilde{a}_n(\gamma)r^{-an}s^{-bn}z^{-n}\Big)e^{\gamma}z^{\partial_\gamma}\\
&= ch(H_+(\gamma, z))ch(S(H_+(\gamma\otimes r^a s^b, z^{-1})^*))e^{\gamma}z^{\partial_\gamma},
\end{aligned}$$

$$\begin{aligned}
&Y^-(\gamma, a, b, z)\\
&= \exp\Big(-\sum_{n\geq 1}\frac{1}{n}\,\widetilde{a}_{-n}(\gamma)r^{an}s^{bn}z^n\Big)\ \exp\Big(\sum_{n\geq 1}\frac{1}{n}\,\widetilde{a}_n(\gamma)z^{-n}\Big)e^{-\gamma}z^{-\partial_\gamma}\\
&= ch(S(H_+(\gamma\otimes r^a s^b, z^{-1})))ch(H_+(\gamma, z)^*)e^{-\gamma}z^{-\partial_\gamma}.
\end{aligned}$$

We note that for $\gamma \in \Gamma^*, l \in \mathbb{Z}$

$$Y^{\pm}(\gamma\otimes r^k s^l, a, b, z) = Y^{\pm}(\gamma, a, b, r^{-k}s^{-l}z). \tag{7.9}$$

It follows from Theorem 7.4 that

$$\begin{aligned}
ch\big(Y^+(\gamma, a, b, z)\big) &= X^+(\gamma, a, b, z)\\
&= \exp\Big(\sum_{n\geq 1}\frac{1}{n}\,a_{-n}(\gamma)z^n\Big)\\
&\qquad\times\exp\Big(-\sum_{n\geq 1}\frac{1}{n}\,a_n(\gamma)r^{-an}s^{-bn}z^{-n}\Big)e^{\gamma}z^{\partial_\gamma}.
\end{aligned}$$

$$\begin{aligned}
ch\big(Y^-(\gamma, a, b, z)\big) &= X^-(\gamma, a, b, z)\\
&= \exp\Big(-\sum_{n\geq 1}\frac{1}{n}\,a_{-n}(\gamma)r^{an}s^{bn}z^n\Big)\\
&\qquad\times\exp\Big(\sum_{n\geq 1}\frac{1}{n}\,a_n(\gamma)z^{-n}\Big)e^{-\gamma}z^{-\partial_\gamma}.
\end{aligned}$$

When $r = s^{-1} = q$ they specialize to the vertex operators $Y^{\pm}(\gamma, k, z)$ studied in [**FJW2**].

Under the new variable (by identifying $a_i(n)$ with $\widetilde{a}_i(n)$) we obtain that

$$\begin{aligned}
&X^+(\gamma_i\otimes s^{-b}, a, b, z)\\
&= \exp\Big(\sum_{n\geq 1}\frac{a_i(-n)}{[n]}s^{-bn}z^n\Big)\ \exp\Big(-\sum_{n\geq 1}\frac{a_i(n)}{[n]}r^{-an}z^{-n}\Big)e^{\gamma}z^{\partial_\gamma},\\
&X^-(\gamma_i\otimes r^{-a}, k, z)\\
&= \exp\Big(-\sum_{n\geq 1}\frac{a_i(-n)}{[n]}s^{bn}z^n\Big)\ \exp\Big(\sum_{n\geq 1}\frac{a_i(n)}{[n]}r^{an}z^{-n}\Big)e^{-\gamma}z^{-\partial_\gamma},
\end{aligned}$$

where $[n]$ is the two-parameter quantum number (4.1).

8. Basic representations and the McKay correspondence

8.1. Two-parameter quantum toroidal algebras. In this subsection we define the two-parameter quantum toroidal algebras$U_{r,s}(\widehat{\widehat{\mathfrak{g}}})$ of simply laced type A, D or E. In particular the two-parameter quantum toroidal algebra contains a special subalgebra -the two-parameter quantum affine algebras $U_{r,s}(\widehat{\mathfrak{g}})$(cf. [**HRZ**]).

Let (A_{ij}) be the two-parameter quantum Cartan $(N+1)\times(N+1)$- Martix(cf. [**HZ, Z**]), For type $A_n^{(1)}$, we have

$$A_{ij} = \begin{pmatrix} rs^{-1} & r^{-1} & 1 & \cdots & 1 & s\\ s & rs^{-1} & r^{-1} & \cdots & 1 & 1\\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots\\ 1 & 1 & 1 & \cdots & rs^{-1} & r^{-1}\\ r^{-1} & 1 & 1 & \cdots & s & rs^{-1}\end{pmatrix}$$

Definition 8.1. The two-parameter quantum toroidal algebra $U_{r,s}(\widehat{\widehat{\mathfrak{g}}})$ is an associative algebra over $\mathbb{K}$ generated by the elements $x_i^{\pm}(k)$, $a_i(m)$, $\omega_i^{\pm1}$, $\omega_i'^{\pm1}$, $\gamma^{\pm\frac{1}{2}}$, $\gamma'^{\,\pm\frac{1}{2}}$, $D^{\pm1}$, $D'^{\,\pm1}$, $(0\leq i\leq N,\ k,\ k'\in\mathbb{Z},\ m,\ m'\in\mathbb{Z}\backslash\{0\})$, subject to the following defining relations:

(D1) $\gamma^{\pm\frac{1}{2}}$, $\gamma'^{\,\pm\frac{1}{2}}$ are central with $\gamma\gamma' = rs$, $\omega_i\,\omega_i^{-1} = \omega_j'\,\omega_j'^{\,-1} = 1$ $(i, j\in I)$, and

$$\begin{aligned}
[\,\omega_i^{\pm1}, \omega_j^{\pm1}\,] &= [\,\omega_i^{\pm1}, D^{\pm1}\,] = [\,\omega_j'^{\,\pm1}, D^{\pm1}\,] = [\,\omega_i^{\pm1}, D'^{\pm1}\,] = 0\\
&= [\,\omega_i^{\pm1}, \omega_j'^{\,\pm1}\,] = [\,\omega_j'^{\,\pm1}, D'^{\pm1}\,] = [D'^{\,\pm1}, D^{\pm1}] = [\,\omega_i'^{\pm1}, \omega_j'^{\,\pm1}\,].
\end{aligned}$$

(D2) $$[\,a_i(m), a_j(m')\,] = \delta_{m+m',0}\frac{(rs)^{\frac{|m|}{2}}(A_{ii}^{\frac{ma_{ij}}{2}} - A_{ii}^{-\frac{ma_{ij}}{2}})}{|m|(r-s)}\cdot\frac{\gamma^{|m|}-\gamma'^{|m|}}{r-s}.$$

(D3) $$[\,a_i(m),\ \omega_j^{\pm1}\,] = [\ a_i(m),\ \omega'^{\pm1}_j\,] = 0.$$

(D4) $$\begin{aligned} D\,x_i^{\pm}(k)\,D^{-1} &= r^k\,x_i^{\pm}(k), & D'\,x_i^{\pm}(k)\,D'^{\,-1} &= s^k\,x_i^{\pm}(k),\\ D\,a_i(m)\,D^{-1} &= r^m\,a_i(m), & D'\,a_i(m)\,D'^{\,-1} &= s^m\,a_i(m).\end{aligned}$$

(D5) $$\omega_i\,x_j^{\pm}(k)\,\omega_i^{-1} = A_{ji}^{\pm1}x_j^{\pm}(k), \qquad \omega_i'\,x_j^{\pm}(k)\,\omega_i'^{\,-1} = A_{ij}^{\mp1}x_j^{\pm}(k).$$

$$
\text{(D6}_1\text{)}\qquad \begin{aligned}[\,a_i(m), x_j^{\pm}(k)\,] &= \pm\tfrac{(rs)^{\frac{|m|}{2}}((rs^{-1})^{\frac{ma_{ij}}{2}}-(rs^{-1})^{-\frac{ma_{ij}}{2}})}{m(r-s)}\\ &\qquad\qquad \cdot\gamma^{\pm\frac{m}{2}}x_j^{\pm}(m{+}k), \qquad \textit{for} \quad m<0,\end{aligned}
$$

$$
\text{(D6}_2\text{)}\qquad \begin{aligned}[\,a_i(m), x_j^{\pm}(k)\,] &= \pm\tfrac{(rs)^{\frac{|m|}{2}}((rs^{-1})^{\frac{ma_{ij}}{2}}-(rs^{-1})^{-\frac{ma_{ij}}{2}})}{m(r-s)}\\ &\qquad\qquad \cdot\gamma'^{\pm\frac{m}{2}}x_j^{\pm}(m{+}k), \qquad \textit{for} \quad m>0,\end{aligned}
$$

$$
\text{(D7)}\qquad \begin{aligned}&x_i^{\pm}(k{+}1)\,x_j^{\pm}(k') - A_{ji}^{\pm1}x_j^{\pm}(k')\,x_i^{\pm}(k{+}1)\\ &= -\Big(A_{ji}A_{ij}^{-1}\Big)^{\pm\frac{1}{2}}\Big(x_j^{\pm}(k'{+}1)\,x_i^{\pm}(k) - A_{ij}^{\pm1}x_i^{\pm}(k)\,x_j^{\pm}(k'{+}1)\Big),\end{aligned}
$$

$$
\text{(D8)}\qquad [\,x_i^{+}(k),\ x_j^{-}(k')\,] = \frac{\delta_{ij}}{r-s}\Big(\gamma'^{-k}\,\gamma^{-\frac{k+k'}{2}}\,\omega_i(k{+}k') - \gamma^{k'}\,\gamma'^{\frac{k+k'}{2}}\,\omega_i'(k{+}k')\Big),
$$

where $\omega_i(m)$, $\omega_i'(-m)$ ($m\in\mathbb{Z}_{\geq 0}$) with $\omega_i(0)=\omega_i$ and $\omega_i'(0)=\omega_i'$ are defined by:

$$
\sum_{m=0}^{\infty}\omega_i(m)z^{-m} = \omega_i\exp\Big((r{-}s)\sum_{\ell=1}^{\infty}a_i(\ell)z^{-\ell}\Big);
$$

$$
\sum_{m=0}^{\infty}\omega_i'(-m)z^{m} = \omega_i'\exp\Big(-(r{-}s)\sum_{\ell=1}^{\infty}a_i(-\ell)z^{\ell}\Big),
$$

with $\omega_i(-m)=0$ and $\omega_i'(m)=0,\ \forall\, m>0$.

$$
\text{(D9}_1\text{)}\qquad x_i^{\pm}(m)x_j^{\pm}(k) = \langle j,i\rangle^{\pm1}x_j^{\pm}(k)x_i^{\pm}(m), \qquad a_{ij}=0,
$$

(D9$_2$)

$$
\begin{aligned}Sym_{m_1,\cdots m_n}\textstyle\sum_{k=0}^{n=1-a_{ij}}(-1)^k(r_is_i)^{\pm\frac{k(k-1)}{2}}\begin{bmatrix}1-a_{ij}\\ k\end{bmatrix}_{\pm i}x_i^{\pm}(m_1)\cdots x_i^{\pm}(m_k)x_j^{\pm}(\ell)\\ \times x_i^{\pm}(m_{k+1})\cdots x_i^{\pm}(m_n)=0, \qquad a_{ij}<0, \quad 0\leq j<i<N,\end{aligned}
$$

(D9$_3$)

$$
\begin{aligned}Sym_{m_1,\cdots m_n}\textstyle\sum_{k=0}^{n=1-a_{ij}}(-1)^k(r_is_i)^{\mp\frac{k(k-1)}{2}}\begin{bmatrix}1-a_{ij}\\ k\end{bmatrix}_{\mp i}x_i^{\pm}(m_1)\cdots x_i^{\pm}(m_k)x_j^{\pm}(\ell)\\ \times x_i^{\pm}(m_{k+1})\cdots x_i^{\pm}(m_n)=0, \qquad a_{ij}<0, \quad 0\leq i<j<N,\end{aligned}
$$

where *Sym* denotes symmetrization with respect to the indices (m_1,m_2),

$$
\begin{bmatrix}n\\ k\end{bmatrix}_{\mp} = \frac{[n]_{\mp}!}{[n-k]_{\mp}![k]_{\mp}!},
$$

and the $\mp$ specifies the substitution of parameters r, s by $r^{\mp}, s^{\mp}$.

The generating functions is defined by:

$$
x_i^{\pm}(z)=\sum_{k\in\mathbb{Z}}x_i^{\pm}(k)z^{-k}, \quad \omega_i(z)=\sum_{m\in\mathbb{Z}_+}\omega_i(m)z^{-m}, \quad \omega_i'(z)=\sum_{n\in-\mathbb{Z}_+}\omega_i'(n)z^{-n}.
$$

REMARK 8.2. Replacing the index $0\leq i\leq N$ by $1\leq i\leq N$ in the above definition, we get the corresponding definition for two-parameter quantum affine algebras (cf. [**HZ, Z**]).

In the case of typr A, the two-parameter quantum toroidal algebra $U_{r,s}(\widehat{\widehat{\mathfrak{g}}})$ admits a further deformation $U_{r,s,\kappa}(\widehat{\widehat{\mathfrak{g}}})$. Let (b_{ij}) be the skew-symmetric$(N+1)\times(N+1)-$ matrix

$$\begin{pmatrix} 0 & 1 & 0 & \cdots & 0 & -1 \\ -1 & 0 & 1 & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & 0 & \cdots & -1 & 0 \end{pmatrix}$$

Definition 8.3. Let κ be an element of $\mathbb{K}^*$. The two-parameter quantum toroidal algebra $U_{r,s,\kappa}(\widehat{\widehat{\mathfrak{g}}})$ is an associative algebra over $\mathbb{K}$ generated by the elements $x_i^{\pm}(k)$, $a_i(m)$, $\omega_i^{\pm 1}$, $\omega_i'^{\pm 1}$, $\gamma^{\pm\frac{1}{2}}$, $\gamma'^{\,\pm\frac{1}{2}}$, $D_1^{\pm 1}$, $D_1'^{\pm 1}$, $D_2^{\pm 1}$, $D_2'^{\pm 1}$ $(0\le i\le N$, k, $k'\in\mathbb{Z}$, m, $m'\in\mathbb{Z}\backslash\{0\}$, $l, l'=1,2)$, subject to the following defining relations:

(T1) $\gamma^{\pm\frac{1}{2}}$, $\gamma'^{\,\pm\frac{1}{2}}$ are central with $\gamma\gamma'=rs$, $\omega_i\,\omega_i^{-1}=\omega_j'\,\omega_j'^{\,-1}=1$ $(i,j\in I)$, and

$$\begin{aligned}[\,\omega_i^{\pm 1},\omega_j^{\pm 1}\,]=[\,\omega_i^{\pm 1},D_l^{\pm 1}\,]=[\,\omega_j'^{\,\pm 1},D_l^{\pm 1}\,]=[\,\omega_i^{\pm 1},D_l'^{\pm 1}\,]=0\\ =[\,\omega_i^{\pm 1},\omega_j'^{\,\pm 1}\,]=[\,\omega_j'^{\,\pm 1},D_l'^{\pm 1}\,]=[D_l'^{\,\pm 1},D_{l'}^{\pm 1}]=[\,\omega_i'^{\pm 1},\omega_j'^{\,\pm 1}\,].\end{aligned}$$

$$\text{(T2)}\qquad [\,a_i(m),a_j(m')\,]=\delta_{m+m',0}\frac{(rs)^{\frac{|m|}{2}}(A_{ii}^{\frac{ma_{ij}}{2}}-A_{ii}^{-\frac{ma_{ij}}{2}})}{|m|(r-s)}\cdot\frac{\gamma^{|m|}-\gamma'^{|m|}}{r-s}\kappa^{mb_{ij}}.$$

$$\text{(T3)}\qquad [\,a_i(m),\ \omega_j^{\pm 1}\,]=[\,a_i(m),\ \omega'^{\pm 1}_j\,]=0.$$

$$\text{(T4)}\qquad \begin{aligned} D_1\,x_i^{\pm}(k)\,D_1^{-1}&=r^k\,x_i^{\pm}(k), & D_1'\,x_i^{\pm}(k)\,D_1'^{\,-1}&=s^k\,x_i^{\pm}(k),\\ D_1\,a_i(m)\,D_1^{-1}&=r^m\,a_i(m), & D_1'\,a_i(m)\,D_1'^{\,-1}&=s^m\,a_i(m),\\ D_2\,x_i^{\pm}(k)\,D_2^{-1}&=r^{\pm\delta_{i0}}\,x_i^{\pm}(k), & D_2'\,x_i^{\pm}(k)\,D_2'^{\,-1}&=s^{\pm\delta_{i0}}\,x_i^{\pm}(k),\\ D_2\,a_i(m)\,D_2^{-1}&=a_i(m), & D_2'\,a_i(m)\,D_2'^{\,-1}&=a_i(m).\end{aligned}$$

$$\text{(T5)}\qquad \omega_i\,x_j^{\pm}(k)\,\omega_i^{-1}=A_{ji}^{\pm 1}x_j^{\pm}(k),\qquad \omega_i'\,x_j^{\pm}(k)\,\omega_i'^{\,-1}=A_{ij}^{\mp 1}x_j^{\pm}(k).$$

$$\text{(T6}_1\text{)}\qquad \begin{aligned}[\,a_i(m),x_j^{\pm}(k)\,]=\pm\tfrac{(rs)^{\frac{|m|}{2}}((rs^{-1})^{\frac{ma_{ij}}{2}}-(rs^{-1})^{-\frac{ma_{ij}}{2}})}{m(r-s)}\\ \cdot\gamma^{\pm\frac{m}{2}}\kappa^{mb_{ij}}x_j^{\pm}(m{+}k),\qquad \textit{for}\quad m<0,\end{aligned}$$

$$\text{(T6}_2\text{)}\qquad \begin{aligned}[\,a_i(m),x_j^{\pm}(k)\,]=\pm\tfrac{(rs)^{\frac{|m|}{2}}((rs^{-1})^{\frac{ma_{ij}}{2}}-(rs^{-1})^{-\frac{ma_{ij}}{2}})}{m(r-s)}\\ \cdot\gamma'^{\pm\frac{m}{2}}\kappa^{mb_{ij}}x_j^{\pm}(m{+}k),\qquad \textit{for}\quad m>0,\end{aligned}$$

$$\text{(T7)}\qquad \begin{aligned}&\Big(\kappa^{b_{ij}}z-(A_{ij}A_{ji})^{\pm\frac{1}{2}}w\Big)\,x_i^{\pm}(z)x_j^{\pm}(w)\\ &\qquad=\Big(\kappa^{b_{ij}}A_{ji}^{\pm 1}z-(A_{ji}A_{ij}^{-1})^{\pm\frac{1}{2}}w\Big)\,x_j^{\pm}(w)\,x_i^{\pm}(z).\end{aligned}$$

$$\text{(T8)}\qquad [\,x_i^{+}(k),\ x_j^{-}(k')\,]=\frac{\delta_{ij}}{r-s}\Big(\gamma'^{-k}\,\gamma^{-\frac{k+k'}{2}}\,\omega_i(k{+}k')-\gamma^{k'}\,\gamma'^{\frac{k+k'}{2}}\,\omega_i'(k{+}k')\Big),$$

where $\omega_i(m)$, $\omega'_i(-m)$ $(m \in \mathbb{Z}_{\geq 0})$ with $\omega_i(0) = \omega_i$ and $\omega'_i(0) = \omega'_i$ are defined by:

$$\sum_{m=0}^{\infty} \omega_i(m) z^{-m} = \omega_i \exp\Big((r-s)\sum_{\ell=1}^{\infty} a_i(\ell) z^{-\ell}\Big);$$

$$\sum_{m=0}^{\infty} \omega'_i(-m) z^{m} = \omega'_i \exp\Big(-(r-s)\sum_{\ell=1}^{\infty} a_i(-\ell) z^{\ell}\Big),$$

with $\omega_i(-m) = 0$ and $\omega'_i(m) = 0,\ \forall\, m > 0$.

$$\text{(T9}_1\text{)} \qquad x_i^{\pm}(m) x_j^{\pm}(k) = \langle j, i\rangle^{\pm 1} x_j^{\pm}(k) x_i^{\pm}(m), \qquad a_{ij} = 0,$$

(T9$_2$)

$$Sym_{m_1,\cdots m_n} \textstyle\sum_{k=0}^{n=1-a_{ij}} (-1)^k (r_i s_i)^{\pm\frac{k(k-1)}{2}} \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_{\pm i} x_i^{\pm}(m_1)\cdots x_i^{\pm}(m_k) x_j^{\pm}(\ell)$$
$$\times x_i^{\pm}(m_{k+1})\cdots x_i^{\pm}(m_n) = 0, \qquad a_{ij} < 0, \quad 0 \leq j < i < N,$$

(T9$_3$)

$$Sym_{m_1,\cdots m_n} \textstyle\sum_{k=0}^{n=1-a_{ij}} (-1)^k (r_i s_i)^{\mp\frac{k(k-1)}{2}} \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_{\mp i} x_i^{\pm}(m_1)\cdots x_i^{\pm}(m_k) x_j^{\pm}(\ell)$$
$$\times x_i^{\pm}(m_{k+1})\cdots x_i^{\pm}(m_n) = 0, \qquad a_{ij} < 0, \quad 0 \leq i < j < N.$$

Sym denotes symmetrization with respect to the indices (m_1, m_2).

Remark 8.4. Assume that $r = s^{-1} = q$, $U_{r,s,\kappa}(\widehat{\widehat{\mathfrak{g}}})$ is the one-parameter quantum toroidal algebras $U_{q,\kappa}(\widehat{\widehat{\mathfrak{g}}})$ (cf. [**GKV, VV**]).

8.2. A new form of McKay correspondence. In this subsection we let Γ to be a finite subgroup of SU_2 and consider two distinguished choices of the class function ξ in $R_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ introduced in Sect. 4.2.

First we consider

$$\xi = \gamma_0 \otimes ((rs^{-1})^{\frac{1}{2}} + (r^{-1}s)^{\frac{1}{2}}) - \pi \otimes 1_{\mathbb{C}^\times},$$

where π is the character of the two-dimensional natural representation of Γ in SU_2.

The Heisenberg algebra in this case has the following relations (cf. Prop. 5.1 and (4.3)).

$$[a_m(\gamma_i), a_n(\gamma_j)] = \begin{cases} m\delta_{m,-n}((rs^{-1})^{\frac{m}{2}} + (r^{-1}s)^{\frac{m}{2}})C, & i = j \\ m\delta_{m,-n} a^1_{ij} C, & i \neq j \end{cases}, \tag{8.1}$$

where a^1_{ij} are the entries of the affine Cartan matrix of ADE type (see (3.4) at $d = 2$).

Recall that the matrix $A^{1,\,1} = (\langle\gamma_i, \gamma_j\rangle^1_\xi) = (a^1_{ij})_{0\leq i,j\leq N}$ is the Cartan matrix for the corresponding affine Lie algebra [**Mc**]. In particular $a^1_{ii} = 2$; $a^1_{ij} = 0$ or -1 when $i \neq j$ and $\Gamma \neq \mathbb{Z}/2\mathbb{Z}$. In the case of $\Gamma = \mathbb{Z}/2\mathbb{Z}$, $a^1_{01} = a^1_{10} = -2$. Let $\mathfrak{g}$ (resp. $\hat{\mathfrak{g}}$) be the corresponding simple Lie algebra (resp. affine Lie algebra) associated to the Cartan matrix $(a^1_{ij})_{1\leq i,j\leq N}$ (resp. A). Note that the lattice $R_{\mathbb{Z}}(\Gamma)$ is even in this case.

We define the normal ordered product of vertex operators as follows.

$$
\begin{aligned}
&: Y^+(\gamma_i, a, b, z) Y^+(\gamma_j, a', b', w) : \\
=&H_+(\gamma_i, z) H(\gamma_j, w) S(H_+(\gamma_i \otimes r^a s^b, z^{-1})^* H_+(\gamma_j \otimes r^{a'} s^{b'}, w^{-1})^*) \\
&\times e^{\gamma_i + \gamma_j} z^{\partial_{\gamma_i}} w^{\partial_{\gamma_j}}, \\
&: Y^+(\gamma_i, a, b, z) Y^-(\gamma_j, a', b', w) : \\
=&H_+(\gamma_i, z) H(-\gamma_j \otimes r^{b'} s^{a'}, w) S(H_+(\gamma_i \otimes r^a s^b, z^{-1})^* H_+(-\gamma_j \otimes r^{a'} s^{b'}, w^{-1})^*) \\
&\times e^{\gamma_i - \gamma_j} z^{\partial_{\gamma_i}} w^{-\partial_{\gamma_j}}.
\end{aligned}
$$

Other normal ordered products are defined similarly.

The identities in the following theorems are understood as usual by means of correlation functions (cf. e.g. [**FJ, J1, HZ, Z**]).

THEOREM 8.5. *Let* $\xi = \gamma_0 \otimes ((rs^{-1})^{\frac{1}{2}} + (r^{-1}s)^{\frac{1}{2}}) - \pi \otimes 1_{\mathbb{C}^\times}$. *Then the vertex operators* $Y^\pm(\gamma_i, a, b, z), Y^\pm(-\gamma_j, a, b, z)$, $\gamma_i \in \Gamma^*, k \in \mathbb{Z}$ *acting on the group theoretically defined Fock space* $\mathcal{F}_{\Gamma \times \mathbb{C}^\times \times \mathbb{C}^\times}$ *satisfy the following relations.*

$$
\begin{aligned}
&Y^+(\gamma_i, a, b, s^{-b} z) Y^+(\gamma_j, a, b, s^{-b} w) \\
&\quad = \epsilon(\gamma_i, \gamma_j) : Y^+(\gamma_i, a, b, s^{-b} z) Y^+(\gamma_j, a, b, s^{-b} w) : \\
&\quad \times \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ (z - r^{-a} s^{-b} w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - r^{-a} s^{-b} (rs^{-1})^{\frac{1}{2}} w)(z - r^{-a} s^{-b} (r^{-1}s)^{\frac{1}{2}} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases},
\end{aligned}
$$

$$
\begin{aligned}
&Y^-(\gamma_i, a, b, r^{-a} z) Y^-(\gamma_j, a, b, r^{-a} w) \\
&\quad = \epsilon(\gamma_i, \gamma_j)^{-1} : Y^-(\gamma_i, a, b, r^{-a} z) Y^-(\gamma_j, a, b, r^{-a} w) : \\
&\quad \times \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ (z - r^{a} s^{b} w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - r^{a} s^{b} (rs^{-1})^{\frac{1}{2}} w)(z - r^{a} s^{b} (r^{-1}s)^{\frac{1}{2}} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases},
\end{aligned}
$$

$$
\begin{aligned}
&Y^+(\gamma_i, a, b, s^{-b} z) Y^-(\gamma_j, a, b, r^{-a} w) \\
&\quad = \epsilon(\gamma_i, \gamma_j) : Y^+(\gamma_i, a, b, s^{-b} z) Y^-(\gamma_j, a, b, r^{-a} w) : \\
&\quad \times \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ (z - r^{-a} s^{b} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - r^{-a} s^{b} (rs^{-1})^{\frac{1}{2}} w)(z - r^{-a} s^{b} (r^{-1}s)^{\frac{1}{2}} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases},
\end{aligned}
$$

$$
\begin{aligned}
&Y^-(\gamma_i, a, b, r^{-a} z) Y^+(\gamma_j, a, b, s^{-b} w) \\
&\quad = \epsilon(\gamma_i, \gamma_j)^{-1} : Y^-(\gamma_i, a, b, r^{-a} z) Y^+(\gamma_j, a, b, s^{-b} w) : \\
&\quad \times \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ (z - r^{a} s^{-b} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - r^{a} s^{-b} (rs^{-1})^{\frac{1}{2}} w)(z - r^{a} s^{-b} (r^{-1}s)^{\frac{1}{2}} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases},
\end{aligned}
$$

$$\begin{aligned}
&Y^+(-\gamma_i,-a,-b,s^{-b}z)Y^+(-\gamma_j,-a,-b,s^{-b}w)\\
&\quad=\epsilon(\gamma_i,\gamma_j):Y^+(-\gamma_i,-a,-b,s^{-b}z)Y^+(-\gamma_j,-a,-b,s^{-b}w):\\
&\quad\times\begin{cases}1 & \langle\gamma_i,\gamma_j\rangle_\xi^1=0\\ (z-r^{-a}s^{-b}w)^{-1} & \langle\gamma_i,\gamma_j\rangle_\xi^1=-1\\ (z-r^{-a}s^{-b}(rs^{-1})^{\frac{1}{2}}w)(z-r^{-a}s^{-b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1=2\end{cases},
\end{aligned}$$

$$\begin{aligned}
&Y^-(-\gamma_i,-a,-b,r^{-a}z)Y^+(-\gamma_j,-a,-b,r^{-a}w)\\
&\quad=\epsilon(\gamma_i,\gamma_j)^{-1}:Y^-(-\gamma_i,-a,-b,r^{-a}z)Y^+(-\gamma_j,-a,-b,r^{-a}w):\\
&\quad\times\begin{cases}1 & \langle\gamma_i,\gamma_j\rangle_\xi^1=0\\ (z-r^{a}s^{b}w)^{-1} & \langle\gamma_i,\gamma_j\rangle_\xi^1=-1\\ (z-r^{a}s^{b}(rs^{-1})^{\frac{1}{2}}w)(z-r^{a}s^{b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1=2\end{cases},
\end{aligned}$$

$$\begin{aligned}
&Y^+(-\gamma_i,-a,-b,s^{-b}z)Y^-(-\gamma_j,-a,-b,r^{-a}w)\\
&\quad=\epsilon(\gamma_i,\gamma_j):Y^+(-\gamma_i,-a,-b,s^{-b}z)Y^-(-\gamma_j,-a,-b,r^{-a}w):\\
&\quad\times\begin{cases}1 & \langle\gamma_i,\gamma_j\rangle_\xi^1=0\\ (z-r^{-a}s^{b}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1=-1\\ (z-r^{-a}s^{b}(rs^{-1})^{\frac{1}{2}}w)(z-r^{-a}s^{b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1=2\end{cases},
\end{aligned}$$

$$\begin{aligned}
&Y^-(-\gamma_i,-a,-b,r^{-a}z)Y^+(-\gamma_j,-a,-b,s^{-b}w)\\
&\quad=\epsilon(\gamma_i,\gamma_j)^{-1}:Y^-(-\gamma_i,-a,-b,r^{-a}z)Y^+(-\gamma_j,-a,-b,s^{-b}w):\\
&\quad\times\begin{cases}1 & \langle\gamma_i,\gamma_j\rangle_\xi^1=0\\ (z-r^{a}s^{-b}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1=-1\\ (z-r^{a}s^{-b}(rs^{-1})^{\frac{1}{2}}w)(z-r^{a}s^{-b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1=2\end{cases},
\end{aligned}$$

REMARK 8.6. Replacing the vertex operator $Y^{\pm}$ by $X^{\pm}$ via the characteristic map *ch* in the above formulas, we get the corresponding formulas for vertex operators $X^{\pm}(\gamma,a,b,z)$ acting on $V_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

Now we consider the second distinguished class function

$$\xi^{r,\,s,\,\kappa}=\gamma_0\otimes((rs^{-1})^{\frac{1}{2}}+(r^{-1}s)^{\frac{1}{2}})-(\gamma_1\otimes\kappa+\gamma_r\otimes\kappa^{-1}),$$

when Γ is a cyclic group of order $N+1$.

In this case the Heisenberg algebra (5.3) has the following relations according to Prop. 5.1:

$$[a_m(\gamma_i),a_n(\gamma_j)]=\begin{cases}m\delta_{m,-n}((rs^{-1})^{\frac{m}{2}}+(r^{-1}s)^{\frac{m}{2}})\kappa^{mb_{ij}}C, & i=j\\ m\delta_{m,-n}a^1_{ij}\kappa^{mb_{ij}}C, & i\neq j\end{cases}, \tag{8.2}$$

where a^1_{ij} are the entries of the affine Cartan matrix of type A and $r\geq 2$. This is the same Heisenberg subalgebra ($c=1$) in $U_{r,s}(\widehat{\mathfrak{g}})$ provided that we identify

$$a_i(n)=\frac{[n]}{n}a_n(\gamma_i).$$

Recall that (b_{ij}) is the skew-symmetric matrix. We need to slightly modify the definition of the middle term in the vertex operators. For each $i = 0, 1, \dots, N$ we define the modified operator $z^{\partial_{\gamma,\kappa}}$ on the group algebra $\mathbb{C}[R_{\mathbb{Z}}(\Gamma)]$ by

$$z^{\partial_{\gamma_i,\kappa}} e^{\beta} = z^{\langle\gamma_i,\beta\rangle_\xi^1} \kappa^{-\frac{1}{2}\sum_{j=1}^{r}\langle\gamma_i,m_j\gamma_j\rangle_\xi^1 b_{ij}} e^{\beta}, \tag{8.3}$$

where $\beta = \sum_j m_j\gamma_j \in R_{\mathbb{Z}}(\Gamma)$.

We then replace the operator $z^{\pm\partial_{\gamma_i}}$ in the definition of the vertex operators $Y^{\pm}(\gamma_i, a, b, z)$ by the operator $z^{\pm\partial_{\gamma_i,\kappa}}$. The formulas in Theorems 7.4 remain true after the term $z^{\pm\partial}$ appearing in the formulas are modified accordingly.

The proof of the following theorem is similar to that of Theorem 8.5.

THEOREM 8.7. *Let* Γ *be a cyclic group of order* $r+1$ *and let* $\xi^{r,\,s,\,\kappa} = \gamma_0 \otimes ((rs^{-1})^{\frac{1}{2}} + (r^{-1}s)^{\frac{1}{2}}) - (\gamma_1 \otimes \kappa + \gamma_r \otimes \kappa^{-1}),$. *The vertex operators* $Y^{\pm}(\gamma_i, a, b, z)$ *and* $Y^{\pm}(-\gamma_i, a, b, z), \gamma_i \in \Gamma^*$ *acting on the group theoretically defined Fock space* $\mathcal{F}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ *satisfy the following relations.*

$$\begin{aligned} &Y^+(\gamma_i, a, b, s^{-b}z)Y^+(\gamma_j, a, b, s^{-b}w)\\ &\quad= \epsilon(\gamma_i,\gamma_j) : Y^+(\gamma_i, a, b, s^{-b}z)Y^+(\gamma_j, a, b, s^{-b}w) :\\ &\quad\times\begin{cases} 1 & \langle\gamma_i,\gamma_j\rangle_\xi^1 = 0\\ \kappa^{-\frac{1}{2}b_{ij}}(z - r^{-a}s^{-b}\kappa^{b_{ij}}w)^{-1} & \langle\gamma_i,\gamma_j\rangle_\xi^1 = -1\\ (z - r^{-a}s^{-b}(rs^{-1})^{\frac{1}{2}}w)(z - r^{-a}s^{-b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1 = 2 \end{cases}, \end{aligned}$$

$$\begin{aligned} &Y^-(\gamma_i, a, b, r^{-a}z)Y^-(\gamma_j, a, b, r^{-a}w)\\ &\quad= \epsilon(\gamma_i,\gamma_j)^{-1} : Y^-(\gamma_i, a, b, r^{-a}z)Y^-(\gamma_j, a, b, r^{-a}w) :\\ &\quad\times\begin{cases} 1 & \langle\gamma_i,\gamma_j\rangle_\xi^1 = 0\\ \kappa^{-\frac{1}{2}b_{ij}}(z - r^{a}s^{b}\kappa^{b_{ij}}w)^{-1} & \langle\gamma_i,\gamma_j\rangle_\xi^1 = -1\\ (z - r^{a}s^{b}(rs^{-1})^{\frac{1}{2}}w)(z - r^{a}s^{b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1 = 2 \end{cases}, \end{aligned}$$

$$\begin{aligned} &Y^+(\gamma_i, a, b, s^{-b}z)Y^-(\gamma_j, a, b, r^{-a}w)\\ &\quad= \epsilon(\gamma_i,\gamma_j) : Y^+(\gamma_i, a, b, s^{-b}z)Y^-(\gamma_j, a, b, r^{-a}w) :\\ &\quad\times\begin{cases} 1 & \langle\gamma_i,\gamma_j\rangle_\xi^1 = 0\\ \kappa^{-\frac{1}{2}b_{ij}}(z - r^{-a}s^{b}\kappa^{b_{ij}}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1 = -1\\ (z - r^{-a}s^{b}(rs^{-1})^{\frac{1}{2}}w)(z - r^{-a}s^{b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1 = 2 \end{cases}, \end{aligned}$$

$$\begin{aligned} &Y^-(\gamma_i, a, b, r^{-a}z)Y^+(\gamma_j, a, b, s^{-b}w)\\ &\quad= \epsilon(\gamma_i,\gamma_j)^{-1} : Y^-(\gamma_i, a, b, r^{-a}z)Y^+(\gamma_j, a, b, s^{-b}w) :\\ &\quad\times\begin{cases} 1 & \langle\gamma_i,\gamma_j\rangle_\xi^1 = 0\\ \kappa^{-\frac{1}{2}b_{ij}}(z - r^{a}s^{-b}\kappa^{b_{ij}}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1 = -1\\ (z - r^{a}s^{-b}(rs^{-1})^{\frac{1}{2}}w)(z - r^{a}s^{-b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_\xi^1 = 2 \end{cases}, \end{aligned}$$

$$
\begin{aligned}
&Y^+(-\gamma_i,-a,-b,s^{-b}z)Y^+(-\gamma_j,-a,-b,s^{-b}w)\\
&\quad=\epsilon(\gamma_i,\gamma_j):Y^+(-\gamma_i,-a,-b,s^{-b}z)Y^+(-\gamma_j,-a,-b,s^{-b}w):\\
&\quad\times\begin{cases}
1 & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=0\\
\kappa^{-\frac{1}{2}b_{ij}}(z-r^{-a}s^{-b}\kappa^{b_{ij}}w)^{-1} & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=-1\\
(z-r^{-a}s^{-b}(rs^{-1})^{\frac{1}{2}}w)(z-r^{-a}s^{-b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=2
\end{cases},
\end{aligned}
$$

$$
\begin{aligned}
&Y^-(-\gamma_i,-a,-b,r^{-a}z)Y^+(-\gamma_j,-a,-b,r^{-a}w)\\
&\quad=\epsilon(\gamma_i,\gamma_j)^{-1}:Y^-(-\gamma_i,-a,-b,r^{-a}z)Y^+(-\gamma_j,-a,-b,r^{-a}w):\\
&\quad\times\begin{cases}
1 & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=0\\
\kappa^{-\frac{1}{2}b_{ij}}(z-r^{a}s^{b}\kappa^{b_{ij}}w)^{-1} & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=-1\\
(z-r^{a}s^{b}(rs^{-1})^{\frac{1}{2}}w)(z-r^{a}s^{b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=2
\end{cases},
\end{aligned}
$$

$$
\begin{aligned}
&Y^+(-\gamma_i,-a,-b,s^{-b}z)Y^-(-\gamma_j,-a,-b,r^{-a}w)\\
&\quad=\epsilon(\gamma_i,\gamma_j):Y^+(-\gamma_i,-a,-b,s^{-b}z)Y^-(-\gamma_j,-a,-b,r^{-a}w):\\
&\quad\times\begin{cases}
1 & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=0\\
\kappa^{-\frac{1}{2}b_{ij}}(z-r^{-a}s^{b}\kappa^{b_{ij}}w) & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=-1\\
(z-r^{-a}s^{b}(rs^{-1})^{\frac{1}{2}}w)(z-r^{-a}s^{b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=2
\end{cases},
\end{aligned}
$$

$$
\begin{aligned}
&Y^-(-\gamma_i,-a,-b,r^{-a}z)Y^+(-\gamma_j,-a,-b,s^{-b}w)\\
&\quad=\epsilon(\gamma_i,\gamma_j)^{-1}:Y^-(-\gamma_i,-a,-b,r^{-a}z)Y^+(-\gamma_j,-a,-b,s^{-b}w):\\
&\quad\times\begin{cases}
1 & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=0\\
\kappa^{-\frac{1}{2}b_{ij}}(z-r^{a}s^{-b}\kappa^{b_{ij}}w) & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=-1\\
(z-r^{a}s^{-b}(rs^{-1})^{\frac{1}{2}}w)(z-r^{a}s^{-b}(r^{-1}s)^{\frac{1}{2}}w) & \langle\gamma_i,\gamma_j\rangle_{\xi}^{1}=2
\end{cases},
\end{aligned}
$$

REMARK 8.8. Replacing the vertex operators $Y^{\pm}$ by $X^{\pm}$ via the characteristic map *ch* we obtain the corresponding results on the space $V_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

8.3. Quantum vertex representations of $U_{r,s}(\widehat{g})$. For each $i=0,\ldots,N$ let

$$\widetilde{a_i}(n)=\frac{[n]}{n}a_n(\gamma_i).$$

It follows from (5.3) and (8.2) that

$$[\widetilde{a_i}(m),\widetilde{a_j}(n)]=\delta_{m,-n}\frac{(rs)^{\frac{m(1-a_{ij})}{2}}[m\langle\gamma_i,\gamma_j\rangle_{\xi}^{1}]}{m}[m]. \tag{8.4}$$

According to McKay, the bilinear form $\langle\gamma_i,\gamma_j\rangle_{\xi}^{1}$ is exactly the same as the invariant form $(\ |\ )$ of the root lattice of the affine Lie algebra $\widehat{\mathfrak{g}}$. This implies that the commutation relations (T2) are exactly the commutation relations (8.4) of the Heisenberg algebra in $U_{r,s}(\widehat{g})$ if we identify $\widetilde{a_i}(n)$ with $a_i(n)$. Thus the Fock space $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ is a level one representation for the Heisenberg subalgebra in $U_{r,s}(\widehat{g})$.

The following theorem gives a q-deformation of the new form of McKay correspondence in [**?**] and provides a direct connection from a finite subgroup Γ of SU_2 to the quantum toroidal algebra $U_{r,s}(\widehat{g})$ of ADE type.

Theorem 8.9. *Given a finite subgroup Γ of SU_2, each of the following correspondence gives a vertex representation of the quantum toroidal algebra $U_{r,\,s}(\widehat{\widehat{g}})$ on the Fock space $\mathcal{F}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$:*

$$\begin{aligned}
x_i^+(n) &\longrightarrow Y_n^+(\gamma_i\otimes s^b, \frac{1}{2}, -\frac{1}{2}),\\
x_i^-(n) &\longrightarrow Y_n^-(\gamma_i\otimes r^a, \frac{1}{2}, -\frac{1}{2}),\\
a_i(m) &\longrightarrow \frac{[m]}{m}a_m(\gamma_i), \qquad \textit{for} \quad m>0;\\
a_i(m) &\longrightarrow \frac{-[-m]}{m}a_m(\gamma_i), \qquad \textit{for} \quad m<0;\\
\gamma &\longrightarrow r, \qquad \gamma' \longrightarrow s;
\end{aligned}$$

or

$$\begin{aligned}
x_i^+(n) &\longrightarrow Y_n^-(-\gamma_i\otimes s^b, -\frac{1}{2}, \frac{1}{2}),\\
x_i^-(n) &\longrightarrow Y_n^+(-\gamma_i\otimes r^a, -\frac{1}{2}, \frac{1}{2}),\\
a_i(m) &\longrightarrow \frac{[m]}{m}a_m(\gamma_i), \qquad \textit{for} \quad m>0;\\
a_i(m) &\longrightarrow \frac{-[-m]}{m}a_m(\gamma_i), \qquad \textit{for} \quad m<0;\\
\gamma &\longrightarrow r, \qquad \gamma' \longrightarrow s;
\end{aligned}$$

where $i=0,\dots,N$, and $m\in\mathbb{Z}$.

Remark 8.10. Replacing $Y^{\pm}$ by $X^{\pm}$ in the above theorem, we obtain a vertex representation of $U_{r,\,s}(\widehat{\widehat{g}})$ in the space $V_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

According to McKay, the bilinear form $\langle\gamma_i,\gamma_j\rangle_\xi^1$ is exactly the same as the invariant form $(\ |\ )$ of the root lattice of the affine Lie algebra $\widehat{\mathfrak{g}}$. Thus the Fock space $S_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ is a level one representation for the Heisenberg subalgebra in $\mathcal{U}_{r,s}(\widehat{\mathfrak{sl}_n})$.

Denote by $\overline{S}_{\Gamma\times\mathbb{C}^\times}$ the symmetric algebra generated by $a_{-n}(\gamma_i)$, $n>0$, $i=1,\dots,N$ over $\mathbb{C}[r^{\pm1},s^{\pm1}]$. $\overline{S}_{\Gamma\times\mathbb{C}^\times}$ is isometric to $\overline{R}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

We define

$$\overline{\mathcal{F}}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}=\overline{R}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}\otimes\mathbb{C}[\overline{R}_{\mathbb{Z}}(\Gamma)]\cong\overline{S}_{\Gamma\times\mathbb{C}^\times}\otimes\mathbb{C}[\overline{R}_{\mathbb{Z}}(\Gamma)].$$

The space $V_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ associated to the lattice $R_{\mathbb{Z}}(\Gamma)$ is isomorphic to the tensor product of the space $\overline{R}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$ and $R_{\mathbb{Z}}(\Gamma)$ as well as the space associated to the rank 1 lattice $\mathbb{Z}\alpha_0$.

The following theorem gives the new form of McKay correspondence in [**FJW2**] for two-parameter case and provides a direct connection from a finite subgroup Γ of SU_2 to the two-parameter quantum affine algebra $\mathcal{U}_{r,s}(\widehat{\mathfrak{sl}_n})$.

Theorem 8.11. *Given a finite subgroup Γ of SU_2, each of the following correspondence gives the basic representation of the two-parameter quantum affine algebra $\mathcal{U}_{r,s}(\widehat{\mathfrak{sl}_n})$ on the Fock space $\overline{\mathcal{F}}_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$:*

$$\begin{aligned}
x_i^+(n) &\longrightarrow Y_n^+(\gamma_i\otimes s^b, \frac{1}{2}, -\frac{1}{2}),\\
x_i^-(n) &\longrightarrow Y_n^-(\gamma_i\otimes r^a, \frac{1}{2}, -\frac{1}{2}),\\
a_i(m) &\longrightarrow \frac{[m]}{m}a_m(\gamma_i), \qquad \mathit{for} \quad m>0;\\
a_i(m) &\longrightarrow \frac{-[-m]}{m}a_m(\gamma_i), \qquad \mathit{for} \quad m<0;\\
\gamma &\longrightarrow r, \qquad \gamma' \longrightarrow s;
\end{aligned}$$

or

$$\begin{aligned}
x_i^+(n) &\longrightarrow Y_n^-(-\gamma_i\otimes s^b, -\frac{1}{2}, \frac{1}{2}),\\
x_i^-(n) &\longrightarrow Y_n^+(-\gamma_i\otimes r^a, -\frac{1}{2}, \frac{1}{2}),\\
a_i(m) &\longrightarrow \frac{[m]}{m}a_m(\gamma_i), \qquad \mathit{for} \quad m>0;\\
a_i(m) &\longrightarrow \frac{-[-m]}{m}a_m(\gamma_i), \qquad \mathit{for} \quad m<0;\\
\gamma &\longrightarrow r, \qquad \gamma' \longrightarrow s;
\end{aligned}$$

where $i=1,\dots,N$.

Remark 8.12. Replacing $Y^\pm$ by $X^\pm$ in the above theorem, we obtain a vertex representation of $\mathcal{U}_{r,s}(\widehat{\mathfrak{sl}_n})$ in the space $V_{\Gamma\times\mathbb{C}^\times\times\mathbb{C}^\times}$.

ACKNOWLEDGMENT

N. Jing would like to thank the support of NSA grant and NSFC Grant (No. 10728102). H. Zhang would like to thank the support of NSFC (No. 10801094) and Shanghai Leading Academic Discipline Project (No. J50101).

References

[A] Abe, E., *Hopf algebras*, Cambridge Tracts in Mathematics, **74**, Cambridge University Press, 1980.

[BGH1] Bergeron, N.; Gao, Y.; Hu, N., *Drinfel'd doubles and Lusztig's symmetries of two-parameter quantum groups*, arXiv.math.RT/0505614, J. Algebra **301** (2006), 378–405.

[BGH2] Bergeron, N.; Gao, Y.; Hu, N., *Representations of two-parameter quantum orthogonal and symplectic groups*, arXiv.math.QA/0510124, AMS/IP Studies in Advanced Mathematics, "Proceedings of the International Conference on Complex Geometry and Related Fields", Vol. **39** (2007), 1–21.

[BH] Bai, X.; Hu, N., *Two-parameter quantum groups of exceptional type E-series and convex PBW-type basis*, arXiv.math.QA/0605179, Algebra Colloquium, (to appear).

[BW1] Benkart, G.; Witherspoon, S., *Two-parameter quantum groups and Drinfel'd doubles*, Alg. Rep. Theory **7** (2004), 261–286.

[BW2] Benkart, G.; Witherspoon, S., *Representatons of two-parameter quantum groups and Schur-Weyl duality*, Hopf algebras, 62–92, Lecture Notes in Pure and Appl. Math., **237**, Dekker, New York, 2004.

[BW3] Benkart, G.; Witherspoon, S., *Restricted two-parameter quantum groups*, Fields Institute Communications, "Representations of Finite Dimensional Algebras and Related Topics in Lie Theory and Geometry", Vol. **40** (2004), 293–318.

[Dr] Drinfel'd, V. G., *A new realization of Yangians and quantized affine algebras*, Soviet Math. Dokl. **36** (1988), 212–216.

[FJ] Frenkel, I.; Jing, N., *Vertex representations of quantum affine algebras*, Proc. Nat'l. Acad. Sci. USA. **85** (1998), 9373–9377.

[FK] Frenkel, I.; Kac, V., *Basic representations of affine Lie algebras and dual resonnace models*, Invent. Math. **62** (1980), 23-66.

[FJW1] Frenkel, I.; Jing, N., Wang, W., *Vertex representations via finite groups and the McKay correspondence*, Int'l. Math. Res. Notice, **4** (2000), 196-222.

[FJW2] Frenkel, I.; Jing, N., Wang, W., *Quantum vertex representations via finite groups and the McKay correspondence*, Commun. Math. Phys., **211** (2000), 365-393.

[GKV] V. Ginzburg, M. Kapranov and E. Vasserot, Langlands reciprocity for algebraic surfaces, Math. Res. Lett. 2 (1995), 147-160.

[HS] Hu, N.; Shi, Q., *The two-parameter quantum group of exceptional type G_2 and Lusztig's symmetries*, arXiv.math.QA/0601444, Pacific J. Math. **230** (2007), 327–345.

[HRZ] Hu, N.; Rosso, M.; Zhang, H., *Two-parameter Quantum Affine Algebra $U_{r,s}(\widehat{\mathfrak{sl}}_n)$, Drinfeld Realization and Quantum Affine Lyndon Basis*, Commun. Math. Phys., **278**, 453-486, (2008)

[HZ] Hu, N.; Zhang, H., *Vertex Representations of Two-parameter Quantum Affine Algebras $U_{r,s}(\widehat{\mathfrak{g}})$: The Simply Laced Cases* , Preprint (2006)

[J1] Jing, N., *Twisted vertex representations of quantum affine algebras*, Invent. Math. **102** (1990), 663–690.

[J2] Jing, N., *On Drinfel'd realization of quantum affine algebras*, Ohio State Univ. Math. Res. Inst. Publ. de Gruyter, Berlin, **7** (1998), 195–206.

[J3] Jing, N., *Quantum Kac-Moody algebras and vertex representations*, Lett. Math. Phys. **44** (1998), 261-271.

[J4] Jing, N., *Vertex representations and McKay correspondence.* Algebra Colloq. **11** (2004), no. 1, 53–70.

[K] Kac, V., *Infinite Dimentional Lie Algebras*, 3rd edition, Cambridge Univ. Press, 1990.

[M] Macdonald, I.G., *Symmetric functions and Hall polynomials*, 2nd ed. Oxford: Clarendon Press, (1995).

[Mc] McKay, J., *Graphs, singularities and finite groups*, Proc. Sympos. Pure Math.,**37**, AMS, (1980), 183-186.

[R] N. Reshetikhin, *Multi-parmater yang-Baxter equation and quantum groups*, Lett. Math. Phys.

[T] Takeuchi, M., *A two-parameter quantization of $GL(n)$,* Proc. Japan Acad. **66** Ser. A (1990), 112–114.

[VV] Varagnolo, V. and Vasserot, E., *Double-loop algebras and the Fock space*, Invent. Math. **133** (1998), 133-159.

[W] Wang, W., *Equivariant K-theory, wreath products and Heisenberg algebra*, Duke Math. J. **103**,(2000), 1–23.

[Z] Zhang, H., *Drinfeld realizations, quantum affine Lyndon bases and vertex representations of two-parameter quantum affine algebras*,Ph. D. thesis, ECNU, Shanghai, China, 2007.

NJ: Department of Mathematics, North Carolina State University, Raleigh, NC 27695, USA
E-mail address: `jing@math.ncsu.edu`

HZ: Department of Mathematics, Shanghai University, Shanghai 200444, China
E-mail address: `hlzhangmath@shu.edu.cn`